How to Cite:

Thamer, S. M., Yahya, M. Q., Mahmood, A. M., & Ahmed, A. G. (2022). Effectiveness of lenalidomide as a topical ointment in mouse models of imiquimod-induced psoriasis. *International Journal of Health Sciences*, *6*(S6), 5477–5489. https://doi.org/10.53730/ijhs.v6nS6.10824

Effectiveness of lenalidomide as a topical ointment in mouse models of imiquimod-induced psoriasis

Sajjad M. Thamer

Ms.c Department of Pharmacology, College of Medicine/ University of Baghdad

Mohammed Qasim Yahya

Assistant Professor, Department of Pharmacology, College of Medicine/ University of Baghdad

Ahmed Mh. Mahmood

Ph.D. Pharmacology, College of Medicine / Al-Nahrain University

Ahmed Ghaith Ahmed

BDS, MSc, Asst. Lecturer, Department of Prosthodontics, Collage of Dentistry/Al-Farahidi University

Abstract---Background: Psoriasis is an inflammatory skin disorder whose cause is unknown. Psoriasis appears to be an immunemediated disease, according to growing evidence. The immunomodulatory effects of lenalidomide inhibit the production of pro-inflammatory cytokines that have been associated with several hematologic cancers. Lenalidomide enhances the host's immune system by regulating T cell proliferation, which causes alterations in inflammation that are associated with the etiology of psoriasis. The purpose of the study: In this trial, the aim was to study the effectiveness of lenalidomide as an ointment in the treatment of psoriasis-induced in mice and to examine the histological differences between the tested groups. Materials and methods: This study was conducted from November 2021 to June 2022. 70 healthy male albino mice, which were randomly divided into seven groups of 10 mice each. Psoriasis was induced by imiquimod in groups (1, 2, 3, 4, 5 and 6). Group 1 received only 5% imiquimod cream, Group 2 received Clobetasol ointment, Group 3 received only the base containing lenalidomide and Groups (4, 5, and 6) received lenalidomide ointment (1%, 2% and 3%, respectively). In Group 7, healthy mice were used as a comparison control. Statistical analysis of the data was performed using SAS (Statistical Analysis System-version 9.1). One-way ANOVA and least significant differences (LSD) post hoc tests were performed

Manuscript submitted: 9 April 2022, Manuscript revised: 27 June 2022, Accepted for publication: 18 July 2022

International Journal of Health Sciences ISSN 2550-6978 E-ISSN 2550-696X © 2022.

to assess significant differences between means. Results: The psoriatic area improved after treatment with lenalidomide. Baker's scoring system explains the histopathological differences between the tested groups. Baker's scoring system (which indicates a combination of nine histological characteristics) reveals significant differences between the groups. Due to its immunomodulatory and anti-inflammatory effects, our results suggest that lenalidomide can treat imiquimod-induced psoriasis in mice. Conclusions: According to the study, different concentrations of lenalidomide can worsen or improve psoriasis in mice. It is effective at a certain dose and has efficacy comparable to that of a standard drug.

Keywords---lenalidomide, immunomodulatory, anti-inflammatory, baker's scoring system, imiquimod-induced psoriasis in mice.

Introduction

Psoriasis is one of the most common, persistent and non-contagious diseases that is characterized by highly proliferative keratinocytes and widespread leukocyte infiltration¹. In North America and Europe, Psoriasis is the most common immune-mediated skin disease in adults². It is characterized by an overactive cellular immune system. It is similar to Crohn's disease, rheumatoid arthritis, multiple sclerosis, and type 1 diabetes in that "A clinical syndrome produced by the activation of T lymphocytes and b lymphocytes, or even both, in the absence of a chronic infection or other identifiable cause," ³. Inflammatory cytokines such as tumor necrosis factor (TNF) are believed to have a major pathogenic role in psoriasis, and other forms of inflammatory leukocytes may also play a role. Infiltrating leucocytes, resident skin cells, and a variety of pro-inflammatory cytokines, chemokines, and chemical mediators released in the skin under the control of the cellular immune system are all associated with psoriasis⁴.

Lenalidomide is a thalidomide analogue that was synthesized by altering the chemical structure of thalidomide to enhance potency and minimize side effects such as side effects of the nervous system (sedation and neuropathy)⁵. Its activity has been shown to be effective as a treatment for a variety of solid and hematological tumors⁵. The Food and Drug Administration (FDA) has approved it for the treatment of patients with myelodysplastic syndromes and multiple myeloma⁶. Lenalidomide has been shown to be an immunomodulator, affecting both the cellular and humoral components of the immune system. Lenalidomide has been shown to modulate cytokine production, T cell co-stimulatory activity, and NK cell cytotoxicity⁶. Lenalidomide has been shown to inhibit the production of pro-inflammatory cytokines TNF-a, IL-1, IL-6, IL-12 and increase the production of anti-inflammatory cytokines IL-10 from human peripheral blood mononuclear cells7. Downregulation of TNF-a secretion is particularly striking and is up to 50,000 times greater compared to thalidomide⁸. TNF-a is a highly pleiotropic cytokine produced primarily by monocytes and macrophages and plays an important role in protective immune responses against bacterial and viral infections. Elevated TNF-a production is involved in the pathogenesis of psoriasis⁹.

The purpose of the study

In this trial, the aim was to study the effectiveness of lenalidomide as an ointment in the treatment of psoriasis-induced in mice and to examine the histological differences between the tested groups.

Materials and Methods

Chemicals and reagents

The chemicals and reagents used in this experiment were as follows: linalidomide powder (Hangzhou hyper chemicals-China), 5% IMQ propionate cream (Meda-Sweden), Clobetasol 17-propionate ointment (GSK), Castor oil (KTC/Germany), petroleum jelly (Battles, Hayward, Bower Ltd/UK), Diethyl Ether (Fluka-Switzerland), Phosphate buffer solution (Pure chemistry–Germany), Distilled water (Iraq) , Paraffin wax (Medite-USA), ethanol99.9%) (Hayman®/UK), Xylene (Scharlau-Spain), Hematoxylin stain, Eosin stain, DPX mounting medium (SyrBio-Switzerland), Scott's tap water (bioGnost®/Croatia), hydrochloric acid (HCl) 1% in 70% ethanol (Riedel-de Han®/Germany). All of the chemicals were of analytical reagent grade.

Preparation of lenalidomide ointment

Linalidomide powder was purchased from Hangzhou Hyper Chemicals-China Limited as white powder and then prepared at the College of Pharmacy, University of Baghdad, as an ointment. The lenalidomide powder is dissolved in castor oil¹², which acts as a levigating agent¹³, and then mixed with petroleum jelly by geometric mixing¹⁴ to make three different concentrations of lenalidomide ointment (1%, 2% and 3%).

Mice and Treatments

70 healthy albino mice, 6-8 weeks old, were purchased from the Iraqi center for cancer research and medical genetics. The Experimental Animal Ethics Committee of the Baghdad Medical College of Medicine approved all procedures, which met national guidelines for the care and use of experimental animals (1697, 6/12/2021). The animals were housed in pathogen-free environments. Water and food are provided ad libitum. The mice are randomly divided into seven groups. Group 1 received Aldara cream (imiquimod 5%) once a day for 6 days on the shaved backs of mice to produce a skin-inflammatory animal model similar to psoriasis (induction group)¹⁶. After imiquimod induction of psoriasis, Group 2 received topically clobetasol ointment once a day for 8 days on the shaved backs of mice. Group 3 received only the base containing lenalidomide (castor oil and petroleum jelly) (free of drugs). Groups 4, 5, and 6 received lenalidomide ointment (1%, 2% and 3%, respectively) to apply topically to the shaved backs of mice once daily for 6 days. In Group 7, only mice without any application (apparently healthy) were used as a comparison control group, as shown in tab.1.

Group	Application	No. mice
G1	Induction	10
G2	standard treated with clobetasol ointment	10
G3	Vehicle treatment (drug-free), placebo treatment	10
G4	Treated with lenalidomide ointment 1%	10
G5	Treated with lenalidomide ointment 2%	10
G6	Treated with 3% lenalidomide ointment	10
G7	Control with no application	10

Table 1 Mice and Treatments

One animal per cage was kept in plastic cages measuring 20x25x35 cm. Before beginning the study procedure, the animals were housed for two weeks in a controlled environment with a temperature of $22 \pm 1^{\circ}$ C and a light schedule of 12-12 hour light/dark cycles, as well as an air vacuum to acclimate to the environment of the animal house. The animals had free access to food, commercial pellets and water¹⁵.

Histological Examination Preparation of the Samples

After 6 days of application of IMQ on the skin for the induction group, the animals were sacrificed humanely by cervical dislocation and the skin treatment region was excised and stored in 10% Phosphate Buffered Saline (PBS) for histopathological examination. On day 14, groups of mice that started treatment on day 7 of induction (day 1 after induction) with lenalidomide, clobetasol and placebo for 8 days were also sacrificed humanely by cervical dislocation, and the skin treatment region was excised and stored in 10% Phosphate Buffered Saline (PBS) for histopathological examination.⁷⁷.

Preparation of formalin-fixed paraffin-embedded tissues¹⁸ (FFPE)

- Tissue Fixation
- Dehydration
- Impregnation
- Embedding tissues in paraffin blocks
- Tissue sectioning and slide preparation
- Hematoxylin and Eosin (H&E) staining of paraffin sections

Assessment of histological analysis of skin tissue

- The light microscope (Genx ®/ USA) was used to capture the zones of a slide that were randomly selected at magnifications of X 10 and X 20.
- Barker's scoring system is used for histopathological examination of wound samples depending on the hematoxylin and eosin staining system to confirm healing as in (Table.2) ¹⁷.

Item		Score
keratin	Munro abscess	2.0
	Hyperkeratosis	0.5
	Parakeratosis	1.0
epidermis	Thining above the papillae	0.5
	Lengthening and Clubbing of rete	1.5
	Ridges	
	Acanthosis	0.5
	Lack of granular layer	1.0
dermis	Lymphocytic infiltrate(mild)	0.5
	Moderate	1.0

Papillary congestion

Marked

Table 2 Baker's scoring system

Statistical analysis

Statistical analysis of the data was performed using SAS (Statistical Analysis System-version 9.1)¹¹. One-way ANOVA and least significant differences (LSD) post hoc tests were performed to assess significant differences between means. P < 0.05 is considered statistically significant¹¹.

Results

After 6 days of IMQ application on the skin on the shaved back of the mice, the charactrisics of psoriasis of round, red papules or plaques with a gray or silvery white dry scale appear on the applied area as shown in (Fig. 1.A). On day 14 (day 8 after induction), the results improved completely with respect to psoriasis after the application of clobetasol (Fig. 1.C) and also completely improved after the application of 1% lenalidomide (Fig. 1.D), but mice treated with clobetasol appeared to lose weight and had secondary skin infections. Mice that were treated with lenalidomide 2% (Fig.1.E) and lenalidomide 3% (Fig.1.F) showed no good improvement, as well as mice treated with placebo showed approximately similar effects to (lenalidomide 2%, 3%) (Fig.1.G). (Fig.1.B) shows a mouse without any applications and appears to be healthy.

2.0

0.5



Figure 1. Various treatments on mice

The results of the skin stained with IMQ showed increased epidermal thickness (hyperkeratosis) and subcutaneous tissue compared to the control group. In addition, abnormal differentiation of keratinocytes with marked parakeratosis (nuclei in the stratum corneum) was observed. Thinning above the papillae, lengthening and clubbing of the rete ridges, acanthosis, and the lack of a granular layer are also observed (Fig. 2.A). In the dermis layer, there is marked lymphocytic infiltrate and papillary congestion (Fig. 2.A). The average score of the induction group was statistically 9.20±0.15a. Clobetasol treatment completely inhibited the increase in thickness. Stratum corneum dropouts were observed in IMQ-treated skin. Although thin above Papillae, the lengthening and clubbing of rete ridges and acanthosis disappear, the granular layer returns. There is no lymphocytic infiltrate or papillary congestion 0.00±0.00d (fig.2.B). Histological examination shows approximately similar results when inducted mice are treated with lenalidomide 1% 0.10±0.10d (fig.2.D). The difference between the clobetasol group and the lenalidomide 1% group was not significantly less than LSD (0.54) as the statistical results show, so the groups show approximately similar results.

The mice group that was treated with 2% lenalidomide showed hyperkeratosis, parakeratosis, lengthening and clubbing of the rete ridges, and acanthosis. There is also lymphocytic infiltrate and papillary congestion 4.25±0.35b (fig.2.E, F). Hyperkeratosis, lengthening and clubbing of rete ridges, and acanthosis were observed in the 3% lenalidomide group. There is also a lymphocytic infiltrate 2.90±0.31c (fig.2.G, H). The group treated with placebo, vehicle of lenalidomide (drug-free) showed lengthening and clubbing of rete ridges and acanthosis. Lymphocytic infiltrate and papillary congestion are found 3.20±0.08b (fig.2.C).

There is a significant difference between the groups (lenalidomide 2%, 3% and placebo groups) and the groups (clobetasol and lenalidomide 1%) groups. As shown in Figure 2.J, no abnormal phenotype was observed in the control groups $0.00\pm0.00d$. These results may provide evidence to suggest that treatment with lenalidomide 1% was effective and approximately similar to standard. Treatment with lenalidomide 2% or lenalidomide 3% did not cause improvement and the characteristics of psoriasis were still present.



Munro abscess, Hyperkeratosis, Parakeratosis, Thinning above papillae,
Lengthening and clubbing of rete ridges, Acanthosis, Lack of granular layer, Papillary congestion, Lymphocytic infiltrate, Keratin,
Epidermis, Dermis

Figure 2. Histological examination of excised skin

Groups	Scores
Group 1 Induction	9.20±0.15a
Group 2 Standard	0.00±0.00d
Group 3 Placebo	3.20±0.08b
Group 4 Lenalidomide 1%	0.10±0.10d
Group 5 Lenalidomide 2%	4.25±0.35b
Group 6 Lenalidomide 3%	2.90±0.31c
Group 7 Normal	0.00±0.00d
LSD	0.54

Table 3 Statistical analysis of Baker's scoring

Means with a different letter are significantly different (P<0.05)



Figure 3. Statistical analysis of Baker's scoring

Discussion

Although the cause of psoriasis is uncertain, some characteristics suggest that it is caused by an immunological response¹. Following skin absorption, Imiquimod can activate monocytes, macrophages, and dendritic cells by binding to Toll-like receptors (TLRs) in plasmacytoid dendritic cells (pDCs) and activating them, leading to massive release of downstream factors such as IFN, IL-17A, IL-22, and TNF in psoriasis lesions and peripheral blood¹⁹. The use of a 5% imiquimod cream generated apparent pathological changes similar to psoriasis in the present investigation. Furthermore, Baker's scores in the imiquimod group were significantly higher than in the control group, indicating progression similar to psoriasis in the imiquimod group. These results revealed that imiquimod therapy can cause and aggravate psoriasis in mice. (Fig.3)

Imiquimod can activate the canonical NF- κ B signaling pathway (Nuclear factor kappa-light-chain-enhancer of activated B cells), which has been linked to the pathogenesis of psoriasis induced by imiquimod²⁰. NF- κ B signaling pathway is responsible for the transcriptional induction of pro-inflammatory cytokines,

chemokines, and additional inflammatory mediators in different types of innate immune cells. These inflammatory mediators can both directly participate in inflammation induction and act indirectly by promoting the differentiation of inflammatory T cells²¹. NF- κ B signaling activates Macrophages rapidly and promotes the secretion of a wide variety of pro-inflammatory cytokines, such as IL-1, IL-6, IL-12, TNF- α and chemokines²¹. NF- κ B also plays a role in the regulation of T cell differentiation and effector function. Upon activation, CD4+ T cells differentiate into different subsets of effector T cells, including Th1, Th2, Th17 and T follicular (Tfh) cells, which secrete different cytokines and mediate different aspects of immune responses, such as IL-17 secretion by Th17 cells, an inflammatory cytokine that recruits monocytes and neutrophils to the site of inflammation in response to invasion by pathogens or self-antigens²¹. Although NF- κ B involved in the generation of Treg cells, Treg cells acquire Th17 inflammatory effector functions under lymphopenic conditions²¹.

T cells are important immune response effectors, and their activation is tightly regulated to avoid auto-reactivity. T cell activation involves the presentation of the peptide fragments displayed by antigen-presenting cells (APCs) to the T cell receptor (TCR), and this interaction gives specificity to the response²². However, this interaction alone is not sufficient if a T cell has to generate an effective response against antigen²². A secondary interaction of B7/CTLA4 on the surface of T cells has a negative signal that regulates T cell activation by inhibiting T cell activation by restricting the availability of B7 for interaction with CD-28²³. CTLA4 has a higher affinity for B7 (20 times) than CD-28. Blocked of this interaction provides the co-stimulatory signal that increases the T cell response and aids in T cell proliferation, differentiation, and survival, followed by a cascade of cytokine and cellular responses²³. In certain situations, T cells or B cells that encounter their antigens in the absence of costimulation can be an ergized or deleted by programmed cell death²⁴. Immunomodulatory drugs (IMiDs), including lenalidomide, act on T cells via the B7/CD28 co-stimulatory pathway by blocking the interaction of B7/CTLA4 Ig²⁵. IMiDs do not up-regulate CD28 and B7 expression in T cells and APCs, respectively, but they can directly induce tyrosine phosphorylation of CD28 in T cells, leading to activation of downstream targets such as PI3K, GRB-2-OS and NF-kb. This might explain their ability to partially overcome CTLA4 Ig blockade^{26,34,35}. T cell costimulation with lenalidomide increases the Th1 cytokine response, leading to increased IFN- and IL-2 secretion, which stimulates clonal T cell proliferation and NK cell activity²⁷.



This model proposes that T-cell activation requires two independent signals. The first is transduced through the T cell receptor (TCR) after engagement by antigen; and the second costimulatory signal is delivered by ligation of a distinct receptor present on the surface of the T cell. This model predicts that the engagement of TCR in the absence of costimulation will not activate T cell²⁸. For decades, topical corticosteroids, particularly high-potency corticosteroids, have been the standard for the treatment of psoriasis. Their success can be linked to a variety of mechanisms of action, including anti-inflammatory, immunosuppressive, and antiproliferative effects. Clobetasol was shown to have a considerable impact. Clobetasol has been shown to have an effect on the IL-23/IL-17A axis in the genesis of psoriasis-like inflammation in mice, reducing the levels of IFN, IL-17A, IL-22, and TNF²⁹. This explains why there is a significant difference between the Clobetasol group and the induction group. (Fig.3), (tab.3)

The results of this study do not show significant differences between the Clobetasol group and the 1% lenalidomide group. The immunomodulatory properties of lenalidomide reduce the generation of pro-inflammatory cytokines, which have been linked to the pathogenesis of psoriasis by altering cytokine production and regulating T cell co-stimulation^{31,32}. Lenalidomide at the lowest concentration of 1% has the lowest or absence of a co-stimulatory signal, leading to T cell death by programing cell death^{24,33,37}. This explains why lenalidomide at 1% concentration has similar results to Clobetasol (Fig.3), (tab.3). As lenalidomide concentrations increased, as in 2% and 3%, co-stimulatory signals increased, leading to T-cell activation. This promotes proliferation and differentiation. As a result, the production of pro-inflammatory cytokines such as TNF and IL-17 increased^{27,34}. (Fig.3), (tab.3)

Conclusions

These findings show that lenalidomide ointment at a certain concentration can improve Imiquimod-induced psoriasis in mice. It is effective and has an efficacy similar to that of a standard drug, with a less thinning effect on the skin by reducing the generation of pro-inflammatory cytokines, which have been linked to the pathogenesis of psoriasis by altering cytokine production and regulating T cell co-stimulation.

References

- 1. Al-Bidri KZ, Salman HA, Al-Hassan Y, Hasan MS. Fibromyalgia Syndrome in a sample of Iraqi patients with psoriasis. Journal of the Faculty of Medicine Baghdad. 2014 ;56(1):49-52.
- 2. Allugunti V.R (2022). A machine learning model for skin disease classification using
- 3. Alzobaidy MA, Alburghaif AH, Alhasany HA, Naji MA. Angiotensin-Converting Enzyme Inhibitors May Increase Risk of Severe COVID-19 Infection. Annals of the Romanian Society for Cell Biology. 2021;25(6):17843–9.
- 4. Al-Zobaidy MJ, Martin W. The ability of asymmetric dimethylarginine (ADMA) or monomethylarginine (L-NMMA) to block endothelium-dependent, nitric oxide-mediated relaxation in rat aorta is inversely related to the efficacy of the relaxant stimulus. Eur J Pharmacol. 2014;741:171-7.
- 5. Araújo FA, Kelmann RG, Araújo BV, Finatto RB, Teixeira HF, Koester LS. Development and characterization of parenteral nanoemulsions containing thalidomide. European Journal of Pharmaceutical Sciences. 2011 Feb 14;42(3):238-45.
- 6. Arrieta J, Cartwright JH, Gouillart E, Piro N, Piro O, Tuval I. Geometric mixing. Philosophical Transactions of the Royal Society A. 2020 Sep 4;378(2179):20200168.
- Bangun, H., Arianto, A. and Rehngenana, E. (2021) 'Anti-ulcer Effect of Gastroretentive Drug Delivery System of Alginate Beads Containing Turmeric Extract Solid Dispersion', Open Access Macedonian Journal of Medical Sciences, 9(A), pp. 19–27.
- 8. Cha IY, Yoo SJ, Jang JH. Recent progress in nanoparticle synthesis via liquid medium sputtering and its applications. Journal of Electrochemical Science and Technology. 2016 Mar 31;7(1):13-26.
- Chanana, M. (2018). Empirical study: relationship between self efficacy and academic performance. International Journal of Health & Medical Sciences, 1(1), 28-34. <u>https://doi.org/10.31295/ijhms.v1n1.36</u>
- 10. Chasset F, Francès C. Current concepts and future approaches in the treatment of cutaneous lupus erythematosus: a comprehensive review. Drugs. 2019 Jul;79(11):1199-215.
- 11. Chen L, Flies DB. Molecular mechanisms of T cell co-stimulation and coinhibition. Nature reviews immunology. 2013 Apr;13(4):227-42.
- a. convolution neural network. International Journal of Computing, Programming and Database Management 3(1), 141-147
- 12. Dellacherie MO, Seo BR, Mooney DJ. Macroscale biomaterials strategies for local immunomodulation. Nature Reviews Materials. 2019 Jun;4(6):379-97.
- 13. D'Souza C, Prince HM, Neeson PJ. Understanding the role of T-cells in the antimyeloma effect of immunomodulatory drugs. Frontiers in Immunology. 2021 Mar 5;12:632399.
- 14. Hafidh RR, Hussein SZ, Malallah MQ, Abdulamir AS. Abu Bakar F. A highthroughput quantitative expression analysis of cancer-related genes in human HepG2 cells in response to limonene, a potential anticancer agent. Current cancer drug targets. 2018;18:807–15.

- 15. Horváth S, Komlódi R, Perkecz A, Pintér E, Gyulai R, Kemény Á. Methodological refinement of Aldara-induced psoriasiform dermatitis model in mice. Scientific reports. 2019 Mar 6;9(1):1-8.
- 16. Hunt DW, Ivanova IA, Dagnino L. DRM02, a novel phosphodiesterase-4 inhibitor with cutaneous anti-inflammatory activity. Tissue barriers. 2020 Jul 2;8(3):1765633.
- 17. Hussein ZA, Shaker NS, Tahseen NJ, Al-Tuhafi AM, Mutee AF. Possible antipsoriasis effect of the methanol extract of Phoenix dactylifera L. seeds. Journal of Herbmed Pharmacology. 2020 Jul 1;9(4):382-90.
- Irrera N, Bitto A, Vaccaro M, Mannino F, Squadrito V, Pallio G, Arcoraci V, Minutoli L, Ieni A, Lentini M, Altavilla D. PDRN, a bioactive natural compound, ameliorates imiquimod-induced psoriasis through NF-κB pathway inhibition and Wnt/β-Catenin signaling modulation. International Journal of Molecular Sciences. 2020 Feb 12;21(4):1215.
- 19. Karim LM, Al-Ani NA, Abbas MH. Psoriasis Treatment by Using Narrowband-UVB Phototherapy. Iraqi Journal of Physics. 2022 ;20(1):57-62.
- 20. Khazem RM, Ibraheem SR. Detection of Some Immunological Parameters in Psoriatic Iraqi Female Patients. Iraqi Journal of Science. 2020:2515-24.
- 21. Klapper W, Klein U, De Silva NS, Zha S, Crowe JL, Dai B, Weilemann A, Okosun J, Fitzgibbon J, Du MQ, Dominguez-Sola D. Malignant Lymphomas: Biology and Molecular Pathogenesis. Walter de Gruyter GmbH & Co KG; 2016 Feb 22.
- 22. Kumar, S. (2022). A quest for sustainium (sustainability Premium): review of sustainable bonds. Academy of Accounting and Financial Studies Journal, Vol. 26, no.2, pp. 1-18
- 23. Kwaśniak K, Czarnik-Kwaśniak J, Maziarz A, Aebisher D, Zielińska K, Karczmarek-Borowska B, Tabarkiewicz J. Scientific reports concerning the impact of interleukin 4, interleukin 10 and transforming growth factor β on cancer cells. Central-European Journal of Immunology. 2019;44(2):190.
- 24. LEVIN EC, SAKO EY, FAMENINI S, WU JJ. SECTION VI Special Considerations. Mild to Moderate and Moderate to Severe Psoriasis (Set). 2019 Jun 24.
- 25. Li Y, Zhang G, Chen M, Tong M, Zhao M, Tang F, Xiao R, Wen H. Rutaecarpine inhibited imiquimod-induced psoriasis-like dermatitis via inhibiting the NF-κB and TLR7 pathways in mice. Biomedicine & Pharmacotherapy. 2019 Jan 1;109:1876-83.
- 26. Liu T, Zhang L, Joo D, Sun SC. NF-κB signaling in inflammation. Signal transduction and targeted therapy. 2017 Jul 14;2(1):1-9.
- 27. Lopalco G, Rigante D, Lopalco A, Emmi G, Venerito V, Vitale A, Capozio G, Denora N, Cantarini L, Iannone F. Safety of systemic treatments for Behçet's syndrome. Expert Opinion on Drug Safety. 2020 Oct 2;19(10):1269-301.
- 28. Nedoszytko B, Sokołowska-Wojdyło M, Ruckemann-Dziurdzińska K, Roszkiewicz J, Nowicki RJ. Chemokines and cytokines network in the pathogenesis of the inflammatory skin diseases: atopic dermatitis, psoriasis and skin mastocytosis. Advances in Dermatology and Allergology/Postępy Dermatologii i Alergologii. 2014 May;31(2):84.
- 29. Obeed RK, Salman HA. Treatment of Chronic Plaque Psoriasis with Etanercept and Methotrexate. Iraq Medical Journal. 2020 26;4(4).

- 30. Ogawa E, Sato Y, Minagawa A, Okuyama R. Pathogenesis of psoriasis and development of treatment. The Journal of dermatology. 2018 Mar;45(3):264-72.
- 31. Owen JA, Punt J, Stranford SA. Kuby immunology. New York: WH Freeman; 2013 Jan 25.
- 32. Paudwal G, Rawat N, Gupta R, Baldi A, Singh G, Gupta PN. Recent advances in solid dispersion technology for efficient delivery of poorly water-soluble drugs. Current Pharmaceutical Design. 2019 Apr 1;25(13):1524-35.
- 33. SAS.2010.SAS/STAT Users Guide for Personal Computer. Release 9.13.SAS Institute, Inc., Cary, N.C., USA.
- 34. Shortt J, Hsu AK, Johnstone RW. Thalidomide-analogue biology: immunological, molecular and epigenetic targets in cancer therapy. Oncogene. 2013 Sep;32(36):4191-202.
- 35. Suryasa, I. W., Rodríguez-Gámez, M., & Koldoris, T. (2021). The COVID-19 pandemic. *International Journal of Health Sciences*, 5(2), vi-ix. https://doi.org/10.53730/ijhs.v5n2.2937
- 36. Tageja N. Lenalidomide-current understanding of mechanistic properties. Anti-Cancer Agents in Medicinal Chemistry (Formerly Current Medicinal Chemistry-Anti-Cancer Agents). 2011 Mar 1;11(3):315-26.
- 37. Thokchom SK, Gulati K, Thakur T, Rai N, Ray A. Dendritic Cells and Immunomodulation: Role in Health and Disease. Current Immunology Reviews. 2017 Aug 1;13(2):132-43.
- 38. Vo MC, Anh-NguyenThi T, Lee HJ, Nguyen-Pham TN, Lakshmi TJ, Jung SH, Kim HJ, Lee JJ. Lenalidomide enhances the function of dendritic cells generated from patients with multiple myeloma. Experimental Hematology. 2017 Feb 1;46:48-55.
- 39. Wang J, Xiang M. Targeting Potassium Channels K v1. 3 and KC a3. 1: Routes to Selective Immunomodulators in Autoimmune Disorder Treatment?. Pharmacotherapy: The Journal of Human Pharmacology and Drug Therapy. 2013 May;33(5):515-28.
- 40. Zalzala HH, Abdullah GA, Abbas MY, Mohammedsalih HR, Mahdi BM. Relationship between human leukocyte antigen DRB1 and psoriasis in Iraqi patients. Saudi Med J. 2018;39(9):886–90.
- 41. Zhang KX, D'Souza S, Upton BA, Kernodle S, Vemaraju S, Nayak G, Gaitonde KD, Holt AL, Linne CD, Smith AN, Petts NT. Violet-light suppression of thermogenesis by opsin 5 hypothalamic neurons. Nature. 2020 Sep;585(7825):420-5.