

How to Cite:

Nataraj, N., Karthik, G., Shantha Kumari, N., & Srinivasa, K. V. (2021). Comparative study of copper, zinc levels in uncontrolled type 2 diabetes mellitus with normal subjects. *International Journal of Health Sciences*, 6(S4), 11748–11753.
<https://doi.org/10.53730/ijhs.v6nS4.11321>

Comparative study of copper, zinc levels in uncontrolled type 2 diabetes mellitus with normal subjects

Dr Nataraj. B

Assistant Professor, Dept of Biochemistry, Dr B R Ambedkar Medical College

Dr Karthik G

Assistant professor, Dept of Biochemistry, Dr B R Ambedkar Medical College

Dr Shantha Kumari N

Associate professor, Dept of Biochemistry, Dr B R Ambedkar Medical College

Dr Srinivasa K V*

Associate professor, Dept of General Medicine, Dr B R Ambedkar Medical College

*Corresponding author

Abstract---Background: The burden of Diabetes is high and increasing across globe and particularly in India. Type 2 Diabetes , accounts for majority of the cases, can lead to multiorgan complications. Direct association of minerals , Trace elements and vitamins in the pathogenesis and natural cause of type 2 diabetes mellitus has been observed in many research studies. Aim: To estimate the levels of Copper , Zinc and HBA1c in uncontrolled Type 2 Diabetes mellitus patient. Material and methods: A cross sectional analytical studies was conducted in DR B R AMC. Study group included 40 known case of uncontrolled Diabetes mellitus. Results: In the present Study serum copper levels in uncontrolled Type 2 Diabetes Mellitus patients was 204.3 ± 49.0 and in controls it was 132.4 ± 20.2 . There was a statistically significant increase in the mean copper values in uncontrolled Type 2 Diabetes Mellitus patients ($P < 0.0001$). Serum Zinc levels in uncontrolled Type 2 Diabetes Mellitus patients was 174.7 ± 31.0 $\mu\text{g}/\text{dl}$ and in the controls it was 207.4 ± 36.6 $\mu\text{g}/\text{dl}$. There was a statistically significant decrease in the mean Zinc values ($P < 0.0001$). Conclusion: Trace elements play a vital role in maintaining the normal metabolism and homeostasis. It is very significant to note that any variation in the normal levels these trace elements has direct link in disturbing the normal metabolic activity very much noted in metabolic disease like Type 2 Diabetes

Mellitus. The role of supplementing the trace elements appropriately may help in better outcome.

Keywords---copper, zinc, HBA1c, uncontrolled type 2 diabetes mellitus.

Introduction

The burden of Diabetes is high and increasing across globe and particularly in India . Estimates in 2019 showed that 77 million had diabetes and expected to rise 134 million by 2045. Type 2 diabetes, accounts for majority of the cases, can lead to multi organ complications. These complications are a significant cause for increased premature morbidity and mortality among individuals with diabetes leading to reduced life expectancy and financial and other costs of diabetes leading to profound economic burden on the Indian health care system Direct association of minerals, trace elements and vitamins in the pathogenesis and natural cause of Type 2 diabetes mellitus has been observed in many research studies. Alteration in the metabolism of several trace elements including copper, zinc have been associated with impaired insulin release , insulin resistance and glucose intolerance.

Aims and objectives of the research were to estimate serum copper , zinc levels in uncontrolled type 2 diabetes mellitus patients and to compare with that of healthy individuals. This study helps to correlate the micro nutrient changes with respect to the uncontrolled Type 2 Diabetes Mellitus Patient in whom there is dynamic derangement happening leading to onset of complications which otherwise will be delayed or prevented in a well controlled Type 2 Diabetes Mellitus Patients and Normal Controls.

Materials and Methods

A cross sectional, analytical study was conducted in Jan 2021 to April 2021 in Dr B R Ambedkar Medical College, Bengaluru. After Ethical committee clearance and Consent of study population we started the study. The study included 40 uncontrolled type 2 Diabetes mellitus patients .The age of the Patient varies from 40 to 60 years. The cases were selected on the basis of simple random sampling method . Metabolic disease altering the levels of Zinc copper were excluded . the results were compared with 40 randomly selected normal healthy individual after obtaining due consent . The controls were age and gender matched . The investigation included serum zinc , copper and HBA1c . The same were compared with controls .

The Investigations included serum copper, zinc and HBA1c . The same were compared with controls. Under all aseptic precautions, about 3ml of venous blood was collected in gel tubes was allowed to clot and it was centrifuged at 4000 rpm for 5 minutes. The serum separated was then used for the estimation of copper, zinc. 2 ml of whole blood was collected separately and was used for estimation of HBA1c Serum copper was estimated by colorimetric method using semi autoanalyser. At PH 4.7 Copper, which is bound to Caeruloplasmin, is released

by a reducing agent. It then reacts with a specific color reagent, 3,5-Di-Br-PAESA-(3,5-Dibromo-2-Pyridylazo)-N-Ethyl-N-(3-Sulphopropyl) aniline, to form a stable, colored chelate. The intensity of the color is directly proportional to the amount of copper in the sample.

Serum Zinc was estimated by colorimetric method with deprotenization by EA-50 semiautoanalyser. Zinc present in the sample is chelated by 5-Br-PAPS [2-(5-bromo-2-Pyridylazo)-5-)N-propyl-N-sulfopropylamino)-phenol in the reagent. The formation of this complex is measured at a wavelength of 560nm. HBA1c is estimated by Latex enhanced Immunoturbidometry method. HBA1c is done as part of categorizing Type 2 Diabetes Mellitus Patients with uncontrolled Blood sugar levels.

Statistical analysis

Data obtained was analysed using statistics software SPSS version 23 . Significance of parameters was tested using Student T test . Results of Parameters are considered significant with P value < 0.005.

Results

The present study analyses the correlation between serum copper and zinc in Uncontrolled Type 2 Diabetes Mellitus. The results were compared with the controls. The study was conducted on 40 uncontrolled Type 2 Diabetes Mellitus patients and 40 healthy controls. The cases and controls were age and sex matched. The age group was between 40-60 years. The mean age in uncontrolled Type 2 Diabetes Mellitus patients was 50.9 ± 5.8 years in controls it was 51.3 ± 4.8 years.

Copper (Cu) : Table showing mean and SD, significant difference in the mean copper (Cu) between the study groups

Between groups	Mean + SD	'P' value	Inference
Uncontrolled Type 2 Diabetes Mellitus	204.3 ± 49.0	< 0.0001	Highly significant
Controls	132.4 ± 20.2		

As represented in the table the mean copper levels in uncontrolled Type 2 Diabetes Mellitus patients was 204.3 ± 49.0 µg/dl and in controls it was 132.4 ± 20.2 µg/dl. There was a statistically significant increase in the mean copper values in uncontrolled Type 2 Diabetes Mellitus patients when compared to the control group (P< 0.0001).

Zinc : Table showing the mean and SD, significant difference in the mean Zinc levels (µg/dl) in the study group.

Groups	Mean + SD	'P' value	Inference
Uncontrolled Type 2 Diabetes Mellitus	174.7 ± 31.0	<0.0001	Highly Significant
Controls	207.4 ± 36.6		

As represented in the table, the mean Zinc levels in uncontrolled Type 2 Diabetes Mellitus patients was 174.7 ± 31.0 $\mu\text{g/dl}$ and in the controls it was 207.4 ± 36.6 $\mu\text{g/dl}$. There was a statistically significant increase in the mean Zinc values in Uncontrolled Type 2 Diabetes Mellitus when compared to the control group ($P < 0.0001$).

HBA1c

Groups	Mean + SD	'P' value	Inference
Uncontrolled Type 2 Diabetes Mellitus	9.4 ± 0.9	<0.0001	Highly Significant
Controls	5.3 ± 0.4		

Discussion

In this study Zinc levels in uncontrolled Diabetes were lower than the control group. Robert M et al noted marked hyperzincuria in diabetic subjects. The increase in urinary zinc excretion in the diabetic subjects may be a contributing factor to the hypozincemia². In our study serum Zinc levels were found to be significantly reduced which can be due to low nutritional intake or increased excretion. Other studies in support of our findings are, Priyanka Mor et al, found significantly low level of Zinc in patients in comparisons to control⁴. Thowya et al, found Zn levels in controlled diabetic patient and uncontrolled diabetic were lower than the healthy group⁵.

Zinc is essential for the correct processing, storage, secretion, and action of insulin in beta (β)-cells. Insulin is stored inside secretory vesicles or granules, where two Zn^{++} ions coordinate six insulin monomers to form the hexameric-structure on which matured insulin crystals are based.¹⁸ It is also known that like, most other chronic disorders, diabetes increases the excretion of minerals.¹⁹ Hyperglycemia in diabetes is usually associated with hyperzincuria and increased urinary loss of Zn^{++} , which is responsible for decreases in total body Zn¹ Zinc has antioxidant properties; thus it can stabilize macromolecules against radical induced oxidation. Hyperglycemia and hyperinsulinemia increases the production of free radicals and there is evidence that lipid peroxidation is increased in type 2 diabetes mellitus patients¹. Zinc supplementation resulted in a significant reduction of plasma total cholesterol, LDL-c and TAG, while increasing HDL-c levels in patients with type-2 diabetes⁸.

In our study Copper levels in uncontrolled Diabetes were Higher than the control group. The increase in Cu ion levels in patients with diabetes mellitus may be attributed to hyperglycaemia that may stimulate glycation and release of copper ions and this accelerates the oxidative stress, so that, Advanced Glycation End products are formed³⁸, that are involved in the pathogenesis of diabetic complications¹ Majority of plasma copper is transported bound to ceruloplasmin (>95%); rest is bound to albumin, transcuprein and copper-amino acid complexes. Ceruloplasmin is an acute phase reactant, has ferro-O₂-oxidoreductase (pro-oxidant) activity directed towards ferrous ion stimulated lipid peroxidation and formation of hydroxyl radical in Fenton reaction.³⁶ Copper is toxic in its unbound form, causes redox imbalance due to it's highly redox active

nature, which leads to activation of stress sensitive intracellular signaling pathways through Haber-Weiss reaction.

Robert M Walter et al observed that diabetic subjects are characterized by increased plasma copper compared with control subjects². Our study is in agreement with study done by Priyanka Mor et al and T E Omer et al who have observed significant increase in serum copper levels in Type 2 Diabetes Mellitus compared to control subjects. Diabetes complications associated with the increased free radical production leading to oxidative damage, and many of the pathological effects of copper overload are consistent with an oxidative damage to membranes or various macromolecules. Cu is involved in oxidation-reduction reactions and has a dominant role in diverse proteins such as cytochrome oxidase at the terminal end of the mitochondria electron transport chain. This divalent cation is also involved in super oxide dismutase activity. Copper has the capacity to form covalent bounds and it takes part in many redox processes. Copper ions are involved in generation of reactive oxygen species through Fenton reaction, having a pro-oxidant action.⁹

Copper ion is well known for cytotoxic effect due to its redox chemistry. Excess of copper aggravate the hyperglycemia which causes glycation of various protein and also displaces copper from copper binding site of protein and thus further aggravates the hypercuperemia, which lead to more oxidative damage to various tissue and organs.⁹

Conclusion

Trace elements play a vital role in maintaining the normal metabolism and homeostasis. It is very significant to note that any variation in the normal levels these trace elements has direct link in disturbing the normal metabolic activity and over time leads to complication and this is very much noted in metabolic disease like Type 2 Diabetes Mellitus. In our study there is significant decrease in serum Zinc levels and correspondingly significant increase in serum copper levels which correlates with uncontrolled blood sugar levels indicating an association between trace elements role in maintaining normal metabolic Homeostasis. Disturbed normal metabolic homeostasis leads to various complications. The role of these trace elements in metabolic disease like Uncontrolled Type 2 Diabetes Mellitus is important to understand better treatment by appropriately correcting the micro nutrient deficiency which may help in better outcome , further interventional studies in this regard can establish the imporatance of these micronutrients and help in better Patient care.

References

1. Abdul hamid zargar, Nissar ahmad shah, Shariq rashid masoodi,et al. Copper, Zinc, and Magnesium levels in non-insulin dependent Diabetes mellitus 1998;74:665-668.
2. Alemayehu toma, Eyasu makonnen, Getnet yimer . Role of zinc in diabetes mellitus, oxidative stress and other human healthy: a review article 2013; 1(11): 411-426.

3. Bibi hajra, Shaukat ali orakzai, Uzma faryal, et al. Insulin sensitivity to trace metals (chromium, manganese) in type 2 Diabetic patients and non Diabetic individuals 2016;28(3): 534-536.
4. Ewelina król, Zbigniew krejpcio, Sławomir michalak et al. Effects of combined dietary chromium(iii) propionate complex and thiamine supplementation on insulin sensitivity, blood biochemical indices, and mineral levels in high-fructose-fed rats 2012; 150:350–359.
5. Jayawardena R, Ranasinghe p, Galappatthy P, et al. Effects of zinc supplementation on diabetes mellitus: a systematic review and meta-analysis 2012; 4:13.
6. Jeff geohas, Anne daly, Vijaya juturu, et al. Chromium picolinate and Biotin combination reduces atherogenic index of plasma in patients with type 2 diabetes mellitus: a placebo-controlled, double-blinded, randomized clinical trial 2007; 333(3): 145–153.
7. Kyria jayanne climaco cruz, Ana raquel soares de oliveira, Dilina do nascimento marreiro. Antioxidant role of zinc in diabetes mellitus 2015 ; 6(2): 333-337.
8. Mahardika, I. M. R., Suyasa, I. G. P. D., Kamaryati, N. P., & Wulandari, S. K. (2021). Health literacy is strongest determinant on self-monitoring blood glucose (SMBG) type 2 DM patients during COVID-19 pandemic at public health centre in Tabanan Regency. *International Journal of Health & Medical Sciences*, 4(3), 288-297. <https://doi.org/10.31295/ijhms.v4n3.1752>
9. Priyanga Ranasinghe, Shehani pigera, Priyadarshani galappatthy, et al. zinc and diabetes mellitus: understanding molecular mechanisms and clinical implications 2015; 23:44
10. Priyanka mor, Anjana kanwar rathore , Vanita sonagra, et al. a Study of serum copper, zinc and magnesium in type 2 Diabetes mellitus with complications and without complications 2020; 13: 1927-1930.
11. Ratnawati, I. G. A. A., Suandayani, N. K. T., & Sutapa, G. N. (2019). The linearity of x-ray devices radiation output and its relationship with patient thickness. *International Journal of Physical Sciences and Engineering*, 3(3), 1–6. <https://doi.org/10.29332/ijpse.v3n3.351>
12. Robert m. walter, jr , Janet y. uriu-hare, Katherine lewis olin, et al. Copper, Zinc, Manganese, and magnesium status and complications of diabetes mellitus 1991; 14 : 1050-1056.
13. Sonny bherwani, Ashok kumar ahirwar, Saumya, et al. effect of Serum copper levels in type 2 diabetes mellitus with nephropathy: a case control study in north Indian population. 2017;5(1), 420-424.
14. Suryasa, I. W., Rodríguez-Gámez, M., & Koldoris, T. (2021). Health and treatment of diabetes mellitus. *International Journal of Health Sciences*, 5(1), i-v. <https://doi.org/10.53730/ijhs.v5n1.2864>
15. Tawfeeq f. r. al-auqbi , et al. Serum copper, zinc and cu/zn ratio in diabetics 2007; 20(3): 419- 425.
16. Thiyam romola devi, Davina hijam, Abhishek dubey, et al. Study of serum zinc and copper levels in type 2 Diabetes Mellitus 2018; 3: 2454-7379.
17. Thowyba elhadi omer, Abdelwahab abdien saeed and Salman taha ahmed elmukashfi. assessment of serum zinc and copper levels among sudanese patients with diabetes mellitus type 2 in khartoum state-sudan 2020; 32(4): 120-125.