

**How to Cite:**

Jawarker, R., Channe, P. P., Goudar, S., Singh, P., Rajesh, D., Thobbi, S. U., & Pandey, P. R. (2022). Arthrocentesis- An minimally invasive method for TMJ disc disorder: An original research. *International Journal of Health Sciences*, 6(S7), 2294–2300.  
<https://doi.org/10.53730/ijhs.v6nS7.11854>

## **Arthrocentesis - An minimally invasive method for TMJ disc disorder: An original research**

**Dr. Rohie Jawarker**

MDS, Consultant Oral and Maxillofacial Surgeon, Allahabad, UP

Email: [rohie90j@gmail.com](mailto:rohie90j@gmail.com)

**Dr. Pallavi P. Channe**

Associate Professor, Dept of Oral Medicine and Radiology, Dr. DY Patil Dental College, Dr.D.Y.Patil Vidyapeeth Pune, MH

Email: [pallavi.channe@dpu.edu.in](mailto:pallavi.channe@dpu.edu.in)

**Dr. Siddarth Goudar**

Assistant Professor, Department of Oral and Maxillofacial Surgery, Gadag Institute of Medical Science, Gadag, Karnataka, India

Corresponding Author: Email: [siddarthgoudar1985@gmail.com](mailto:siddarthgoudar1985@gmail.com)

**Dr. Pranendra Singh**

Consultant Maxillo-Facial Surgeon, Prarthna Hospital New Delhi

Email: [rajsngn054@gmail.com](mailto:rajsngn054@gmail.com)

**Dr. Damarasingu Rajesh**

OMFS, PhD Scholar, Dept of OMFS, Narsinhbhai Patel Dental College and Hospital, Sankalchand Patel University, Visnagar, Gujarat

Email: [rajeshoralsurgeon@gmail.com](mailto:rajeshoralsurgeon@gmail.com)

**Dr. Sampada Umesh Thobbi**

Assistant professor, Department of Prosthodontics, Dr.D.Y.Patil Dental College and hospital, pimpri, pune, Dr.D.Y.Patil Vidyapeeth,pune, Maharashtra,India

Email: [sampada99thobbi9@gmail.com](mailto:sampada99thobbi9@gmail.com)

**Dr. Pritee Rajkumar Pandey**

MDS, OMFS, Consultant Oral and MaxilloFacial Surgeon, Mahalaxmi Hospital, Nerul, Navi Mumbai

Email: [Priteepandey1990@gmail.com](mailto:Priteepandey1990@gmail.com)

**Abstract**--TMJ dysfunction is a therapeutic challenge in the oral and maxillofacial clinic. Although TMJ pain and dysfunction can be caused by many different aetiologic factors, the role played by inflammation as an underlying mechanism of pain and dysfunction of the TMJ has played a major role. TMD patients having pain and

tenderness for prolonged period of time will show signs of inflammation biochemically as well as radiographically. Therefore the aim of this study was to evaluate the efficacy of arthrocentesis in patients suffering from TMJ disc disorders. Materials and Methods A Total Of 60 Subjects Suffering From Tmj Disc Disorders Were Selected And Treated By Arthrocentesis. The Subjects Were Followed Up For A Period Of Two Year. Results The mean maximal mouth opening prior to arthro-centesis was 30.10 mm and after the procedure the mean maximal mouth opening was 45.3 mm. The mean right and left lateral movements before arthrocentesis were 6.15mm and 6.59mm respectively, and the mean right and left lateral movements of 9.09 and 9.21 respectively were present after the procedure. The mean degree of pain before arthrocentesis was 7.7, and after the procedure the mean degree of pain was 1.03 as per the visual analogue scale. Conclusion The findings of this study suggested potential utility of arthrocentesis in the management of TMJ disc disorders.

**Keywords**---TMJ pain, Maximal mouth opening, Lateral mandibular movements.

## **Introduction**

Temporomandibular joint (TMJ) disorders accounts for about 10% of the population with female predilection<sup>1</sup>. TMJ dysfunction is a therapeutic challenge in the oral and maxillofacial clinic. Although TMJ pain and dysfunction can be caused by many different aetiologic factors, the role played by inflammation as an underlying mechanism of pain and dysfunction of the TMJ has played a major role. TMD patients having pain and tenderness for prolonged period of time will show signs of inflammation biochemically as well as radiographically<sup>2</sup>. Various treatment modalities available for TMD patients are, arthrocentesis, arthroscopic lysis and lavage and arthrotomy<sup>3</sup>. An important change in the therapeutic approach occurred with the introduction of arthroscopic lavage and lysis for the treatment of TMDs. The success of arthroscopy has led to the use of arthrocentesis as a simple therapeutic modality with a satisfactory outcome<sup>4</sup>. The procedure of arthrocentesis involves aspiration of the joint cavity with injection of therapeutic substance in the superior joint space that causes release of the disc and ultimately resulting in increased mouth opening<sup>5-7</sup>. Arthrocentesis has proved to be a minimally invasive treatment modality, relatively safe, repeatable and it can be done on out patients under local anaesthesia which significantly reverts the mouth opening to a normal range. It is an effective method of normal disc-condyle-fossa relationship<sup>8</sup>. Several authors have conducted studies to detect effectiveness of arthrocentesis in various TMJ disc disorders. Hence, the present study evaluated the effectiveness of arthrocentesis in TMJ disc disorder patients.

## **Material and Methods**

In this study, the minimum of 60 patients with TMJ disc disorder who were clinically diagnosed as per the norms laied down by Kaplan were selected<sup>7</sup>. In all, extreme care was taken to selectively. The clinical examination was done

including the evaluation of the maximal mouth opening which was measured by the distance between the incisal edges of the upper and lower incisors with the help of a caliper. Determination of the 'range of the lateral and protrusive mandibular movement' measured by the distance between the upper and lower midline on lateral and forward movements by using a caliper. Pain level and location were determined by the patients self assessment using VAS ranging from 0 to 108. After thorough TMJ evaluation and clinical examination, orthopantomograph and transpharyngeal view of the patient were taken. The diagnosis of TMJ disc disorder was made on the basis of history, clinical and radiologic examination as per the norms laied down by Kaplan<sup>7</sup>. After proper preparation of the target site, external auditory meatus were blocked with cotton soaked in normal saline. Two points of needle insertion were marked over the skin of the affected joint indicating the articular fossa and eminence. A line was drawn from the middle of the tragus to the outer canthus. The posterior entrance point was located along the canthotragal line, 10mm from the middle of the tragus and 2mm below the line. This posterior point is only for pumping the fluid into the upper compartment to increase the hydraulic pressure within the joint. The anterior point of entry was placed 10mm farther along the line and 10mm below it. This was followed by the injection of local anaesthesia to block the auriculotemporal nerve. An 18-gauge needle was then inserted into the superior compartment of the anterior auricular fossa (posterior mark), followed by the injection of 2-3ml of Ringer's Lactate solution to distend the joint space. Then 18 gauge needle was inserted for the fluid to come out from the superior compartment. Ringer's Lactate solution was then connected to one of the needle with sufficient pressure to assure the free flow of 200ml solution during 15-20min period which was achieved by inserting a syringe at an elevation of 1cm above the level of the joint. During the procedure, exact timing of re-establishment of normal mouth opening determined by having the patient to make the repeated attempts to open the mouth. On termination of the procedure steroid (dexamethasone 8mg) was injected into the joint space followed by the removal of needle. Postoperatively antibiotic and analgesics with muscle relaxants for 2 weeks were prescribed. Follow up of the patient was done every 6 months for 2 years.

## Results

The mean maximal mouth opening prior to arthro-centesis was 30.10 mm and after the procedure the mean maximal mouth opening was 45.3 mm (Table 1). The mean right and left lateral movements before arthrocentesis were 6.15mm and 6.59mm respectively, and the mean right and left lateral movements of 9.09 and 9.21 respectively were present after the procedure (Table 2). The mean degree of pain before arthrocentesis was 7.7, and after the procedure the mean degree of pain was 1.03 as per the visual analogue scale. The maximal mouth opening prior to arthrocentesis ranged from 20-44mm with a mean value of 30.10mm and a SD of 9.86mm. The maximal mouth opening 2 year following arthrocentesis ranged from 41-50mm with a mean value of 45.3mm and a SD of 2.56mm. The right lateral movement prior to arthrocentesis ranged from 5-10mm with a mean value of 7.05mm and a SD of 1.25mm. The right lateral movement 2 year following arthrocentesis ranged from 8-10 mm with a mean value of 9.21mm and a SD of 0.61mm. The left lateral movement prior to arthrocentesis ranged from 5.5-10mm

with a mean value of 7.19mm and a SD of 1.26mm. The left lateral movement following arthrocentesis ranged from 8-11mm with a mean value of 9.11mm and a SD of 0.70mm. The degree of pain experienced by patients before arthrocentesis ranged from 6-10 with a mean value of 7.7 and a SD of 1.1. The degree of pain after arthrocentesis ranged from 1-4 with a mean value of 1.03 and a SD of 1.16.

## Discussion

Temporomandibular disorders which will present itself as pain, clicking sound and deviation are collectively the disorders of joint and muscles<sup>9</sup>. It has been suggested that classification, diagnosis and treatment of TMJ pain and dysfunction can be based on the position and shape of the articular disc. The question of whether disc displacement is the result, cause, or an accompanying factor in dysfunction remains open to debate<sup>10</sup>. The management of refractory pain in the TMJ is both challenging and controversial for Oral and Maxillofacial Surgeons. Controversy continues to surround the role of surgery in the management of pain and dysfunction of the TMJ, although only about 5% of all patients being treated for TMJ disorders are actually operated<sup>11</sup>. The success rate of non-surgical treatments according to many authors is approximately 60%. However, certain group of patients receiving non-surgical treatment may remain unresponsive, thereby prolonging their suffering and treatment dissatisfaction<sup>12</sup>. As the procedures like Arthrocentesis and Arthroscopy are having high success rates with minimal complication, they are used most commonly as a first line of surgical treatment along with nonsurgical treatment. Arthrocentesis is minimally invasive technique with high success rates, very easy to perform and very effective in relieving the pain and ultimately increasing the mouth opening<sup>13</sup>. Thus with the aim to know the efficacy of arthrocentesis in patients suffering from TMJ disorders, the present study was conducted. The present study showed significant increase in the mouth opening from 30.03mm to 45.3mm. The findings were similar to those of Nitzan DW<sup>14</sup> et al., (24.1mm to 42.7mm), Dimitroulis G<sup>15</sup> et al., (24.6mm to 42.3mm), Fridrich KL<sup>16</sup> et al., (33mm to 41mm), Hosaka H<sup>17</sup> et al., (30.6mm to 44.5mm), Nitzan DW<sup>14</sup> et al., (23.1mm to 44.26mm), Carvajal WA<sup>18</sup> et al., (25.3mm to 43.8mm), Nitzan DW<sup>19</sup> et al., (24.4mm to 43.20mm), and Dhaif G<sup>6</sup> et al., (13.1mm to 43.7mm). The findings were higher to those of Yeung RWK<sup>20</sup> et al., (38.2mm to 39.8mm), and Onder ME<sup>21</sup> et al., (33.6mm to 38mm). The variations in the values could be due to the differences in the use of intra-articular medications used for the treatment as is the case in the study done by Yeung RWK et al., and Onder ME et al., who used hyaluronic acid for intra-articular injection. But in our study we have used Ringer's Lactate solution for intra-articular injection<sup>21</sup>. There was significant increase in the right and left lateral movement. The values were in consistent with that of Nitzan DW<sup>14</sup> et al., (6.30mm to 9.40mm), who used intra articular injection of steroids as was done in our study, however the values were slightly higher to those of Nitzan DW<sup>14</sup> et al., (4.8mm to 8.2mm) and the values were slightly lower compared to the study done by Nitzan DW<sup>19</sup> et al., (3.75mm to 10.5mm) who did not use intra-articular steroid. The variations in the values could be due to the differences in the use of intra-articular medications used for the treatment as is the case in the study done by Nitzan D W et al., who did not use intra-articular steroids<sup>14</sup>. The degree of pain experienced by patients before arthrocentesis ranged from 6-10 with a mean value of 7.7 and a SD of 1.1. The degree of pain after arthrocentesis ranged

from 1-4 with a mean value of 1.03 and a SD of 1.16. The probable explanation that could be given due to the variations in the values is short follow up period and variations in the postoperative medications given. In our study, we have used two needle technique along with injection of dexamethasone as described by Nitzan DW<sup>14</sup> et al., and postoperatively antibiotics and analgesics with muscle relaxants were prescribed for two weeks. In the study done by Dimitroulis G<sup>15</sup> et al., and Nitzan DW<sup>19</sup> et al., intra articular injection of steroid was not given. In another study done by Hosaka H<sup>17</sup> et al., single needle technique was used. Postoperative medication of Diazepam 10mg was used in the study done by Dhaif G<sup>6</sup> et al. Intra articular injection of hyaluronic acid was given and there was no significant increase in the mouth opening and reduction in the intensity of pain. The use of intra articular steroid as done by, Goutdot<sup>22</sup> et al and Emshoff R<sup>23</sup> and in our study yielded a highly significant results.

### **Conclusion**

Therefore this study revealed significant increase in maximal mouth opening, lateral movements and significant reduction in TMJ pain after a period of 2 year follow up and suggested potential utility of arthrocentesis in the management of TMJ disc disorders.

### **References**

- [1] Ethunandan M. Temporomandibular joint arthrocentesis – more questions than answers? *J Oral Maxillofac Surg.* 2006;64:952–55. [PubMed] [Google Scholar]
- [2] Kopp S, Wenneberg B, Haraldson T, Carlsson GE. The short- term effect of intra-articular injections of sodium hyaluronate and corticosteroid on temporomandibular joint pain and dysfunction. *J Oral Maxillofac Surg.* 1985;43:429–35. [PubMed] [Google Scholar]
- [3] Emshoff R, Rudisch A. Temporomandibular joint internal derangement and osteoarthritis: are effusion and bone marrow edema prognostic indicators for arthrocentesis and hydraulic distension? *J Oral Maxillofac Surg.* 2007;65:66–73. [PubMed] [Google Scholar]
- [4] Nitzan DW, Price A. The use of arthrocentesis for the treatment of osteoarthritic temporomandibular joints. *J Oral Maxillofac Surg.* 2001;59:1154–59. [PubMed] [Google Scholar]
- [5] Nitzan DW. Arthrocentesis – Incentives for using this minimally invasive approach for temporomandibular disorders. *Oral and Maxillofacial Surg Clin N A.* 2006;18:311–28. [PubMed] [Google Scholar]
- [6] Dhaif G, Ali T. TMJ Arthrocentesis for acute closed lock: retrospective analysis of 40 consecutive cases. *Saudi Dental Journal.* 2001;3. [Google Scholar]
- [7] Kaplan A.S., Assael L.A., editors. *Temporomandibular disorders: diagnosis and treatment.* Philadelphia: W.B. Saunders Company; 1991. pp. 663–679. [Google Scholar]
- [8] Smolka W, Iizuka T. Arthroscopic lysis and lavage in different stages of internal derangement of temporomandibular joint: correlation of preoperative staging to arthroscopic findings and treatment outcome. *J Oral Maxillofac Surg.* 2005;63:471–78. [PubMed] [Google Scholar]

- [9] Emshoff R, Rudisch A, Bosch R, Gabner R. Effect of arthrocentesis and hydraulic distension on the temporomandibular joint disk position. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod.* 2000;89:271–77. [PubMed] [Google Scholar]
- [10] Kunjur J, Anand R, Brennan PA, Ilankovan V. An audit of 405 temporomandibular joint arthrocentesis with intra-articular morphine infusion. *Br J Oral Maxillofac Surg.* 2003;41:29–31. [PubMed] [Google Scholar]
- [11] Schmitter M, Zahran M, Duc JMP, Henschel V, Rammelsberg P. Conservative therapy in patients with anterior disc displacement without reduction using 2 common splints: a randomized clinical trial. *J Oral Maxillofac Surg.* 2005;63:1295–303. [PubMed] [Google Scholar]
- [12] Lee SH, Yoon HJ. MRI findings of patients with temporomandibular joint internal derangement: before and after performance of arthrocentesis and stabilization splints. *J Oral Maxillofac Surg.* 2009;67:314–17. [PubMed] [Google Scholar]
- [13] Yura S, Totsuka Y, Yoshikawa T, Inoue N. Can Arthrocentesis release intracapsular adhesions? Arthroscopic findings before and after irrigation under sufficient hydraulic pressure. *J Oral Maxillofac Surg.* 2001;87:874–77. [PubMed] [Google Scholar]
- [14] Nitzan DW, Dolwick MF, Martinez GA. Temporomandibular Joint arthrocentesis: a simplified treatment for severe, limited mouth opening. *J Oral Maxillofac Surg.* 1991;49:1163–67. [PubMed] [Google Scholar]
- [15] Dimitroulis G, Dolwick MF, Martinez A. Temporomandibular joint arthrocentesis and lavage for the treatment of closed lock: a follow-up study. *Br J Oral Maxillofac Surg.* 1995;33:23–27. [PubMed] [Google Scholar]
- [16] Fridrich KL, Wise JM, Zeitler DL. Prospective comparison of arthroscopy and arthrocentesis for temporomandibular joint disorders. *J Oral Maxillofac Surg.* 1996;54:816–20. [PubMed] [Google Scholar]
- [17] Hosaka H, Murakami K, Goto K, Iizuka T. Outcome of arthrocentesis for temporomandibular joint with closed lock at 3 years follow up. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod.* 1996;82:501–04. [PubMed] [Google Scholar]
- [18] Carvajal WA, Laskin DM. Long-Term evaluation of arthrocentesis for the treatment of internal derangements of the temporomandibular joint. *J Oral Maxillofac Surg.* 2000;58:852–55. [PubMed] [Google Scholar]
- [19] Nitzan DW, Samson B, Better H. Long-term outcome of arthrocentesis for sudden-onset, persistent, severe closed lock of the temporomandibular joint. *J Oral Maxillofac Surg.* 1997;55:151–57. [PubMed] [Google Scholar]
- [20] Yeung RWK, Chow RLK, Samman N, Chiu K. Short-term therapeutic outcome of intra-articular high molecular weight hyaluronic acid injection for nonreducing disc displacement of the temporomandibular joint. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod.* 2006;102:453–61. [PubMed] [Google Scholar]
- [21] Onder ME, Tuz HH, Kocyigit D, Kisnisci RS. Long - term results of arthrocentesis in degenerative temporomandibular disorders. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod.* 2009;107:e1–5. [PubMed] [Google Scholar]

- [22] Suryasa, I. W., Rodríguez-Gámez, M., & Koldoris, T. (2021). Get vaccinated when it is your turn and follow the local guidelines. *International Journal of Health Sciences*, 5(3), x-xv. <https://doi.org/10.53730/ijhs.v5n3.2938>
- [23] Goudot P, Jaquinet AR, Hugonnet S, Haefliger W, Richert M. Improvement of pain and function after arthroscopy and arthrocentesis of the temporomandibular joint: a comparative study. *Journal of Cranio-maxillofacial Surgery*. 2000;28:39–43. [PubMed] [Google Scholar]
- [24] Emshoff R, Rudisch A, Bosch R, Strobl H. Prognostic indicators of the outcome of arthrocentesis: A short-term follow-up study. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod*. 2003;96:12–18. [PubMed] [Google Scholar]

Table 1  
Correlation of maximal mouth opening before and after arthrocentesis

Groups	n	Maximal mouth opening	
		Range	Mean±SD
Before the procedure	60	20-44	30.03±9.86
1 year after the procedure	60	41-50	45.3±2.56
t	8.05		
p*	<0.001 HS		

Table 2  
Correlation of lateral movements before and after arthrocentesis

Groups	n	Lateral movements			
		Right		Left	
		Range	Mean±SD	Range	Mean±SD
Before the procedure	60	5-10	7.05±1.25	5.5-10	7.19±1.26
2 year after the procedure	60	8-10	9.21±0.61	8-11	9.11±0.70
t	9.55		6.217		
p*	<0.001 HS		<0.001 HS		