How to Cite:

Mohammad, S. F. (2022). Impact on ruminant performance of feed additives: Review. *International Journal of Health Sciences*, 6(S6), 10066–10077. https://doi.org/10.53730/ijhs.v6nS6.12629

Impact on ruminant performance of feed additives: Review

Sundos Farook Mohammad

Department of Animal Production, College of Agriculture, University of Baghdad, Iraq

*Corresponding author email: sundos.f@coagri.uobaghdad.edu.iq

Abstract---In order to elicit the desired reaction at animals, feed additives must be included in high concentrations in the diet. Microorganism-based feed additives for ruminants have been more popular over the last two decades, and this trend is expected to continue. Dairy farmers may utilize feed additives to enhance the nutritional content of their dairy cows' meals and boost revenues by using them properly. The use of supplements should not be seen as a substitute for well-balanced diets and sound feeding procedures, but rather as an addition to them. Although yeast biomass, a byproduct of the alcoholic beverage industry, has previously been employed as a protein source, research into feeding yeast as a probiotic began in the 1950s. Probiotics like yeast were only employed as a feed additive at a lesser level. It was in the 1940s and 1950s when Saccharomyces cerevisiae (Sc) was first used as a probiotic in tiny doses added to animal feed. Anaerobic cellulitis bacteria count and ruminal PH value may be improved by adding yeast to ruminant diets, according to research. mineral absorption is influenced by the patterns of volatile fatty acids.

Keywords---saccharomyces cerevisiae, ruminants, performance in yeast.

Introduction

Fermentation in the rumen may be improved by using feed additives, which in turn help the animals absorb nutrients more efficiently. Microbes such Direct-Feed Microbial (DFM), yeast cultures, and live yeast cultures (LYC) were addressed by Denev et.al.(2007). Animal feed microbes were also included on the list in a well-written essay, (Fuller ,1989) summarizes some of the most often held beliefs regarding the benefits of DFM. Ewe gain in weight and feed conversion ratio are aided by both bacteria found in rumen(Khatta,et.al.2020). VFA levels were increased, pH was maintained, and lactate levels were lowered when yeast

International Journal of Health Sciences ISSN 2550-6978 E-ISSN 2550-696X © 2022.

Manuscript submitted: 9 May 2022, Manuscript revised: 18 July 2022, Accepted for publication: 27 August 2022 10066

was added to a rat diet. An increase in feed utilization and enhanced rumen fermentation may also help raise milk production and animal performance (Amin and Mao,2021). However, according to Bayat et al. (2015), DFM supplementation had no influence on the growth rate of finishing cattle. As a probiotic, Saccharomyces cerevisiae (Sc) has been shown to improve the health of calves fed 10g/ calf/day of yeast culture Maamouri and Ben Salem (2021).

Natural Supplements for Animal Feed

Animals may benefit greatly from the use of natural feed additives such as medicinal herbs in their diets. With the addition of medicinal herbs like Nigella sativa and rosemary to limited concentration diets, the weight growth of lamb was significantly increased (Hassan and Hassan ,2010a). When given a diet enriched with Nigella Sativa (Ns), Karadzic sheep gained more weight than those fed diets lacking Ns (Hassan and Hassan,2009a). Serum urea nitrogen (SUN) was considerably (P 0.05) greater in lambs given rosemary-infused diets, according to Hassan and Hassan (2009b). There is some evidence to suggest that herbal feed additives may enhance gastrointestinal function in vitro, according to Hua-wei and Jian-ming (2011). That the performance of the lambs might be enhanced by using the herbal growth enhancer. Microbes in the rumen are raised when medicinal herbs are used as feed additives to influence the microbes (Ruiz Garcia, et.al.2011).

Additives made from microbes

Various microbial feed additives are used into ruminant diets for a variety of reasons. It has been shown in research that the usage of probiotics in nonruminants, such as calves and lambs, may help them gain weight (Mirzaei and Hari ,2012). In addition to humans, ruminants also benefit from the usage of probiotics. Dry matter (DM), organic matter (OM), metaboilizable energy (ME), and total intake (TN) in lambs given Iraqi probiotic supplements was significantly increased when S. cerevisiae (Sc) and Aspergillus or fungal (Salim, et.al. 2009) fermentation extracts were supplied. But according to Seo et al. (2012), lactic acid bacteria (LAB) and other microorganisms, such as Lactobacillus, Bifidobacterium, enterococcus, streptococcus, bacteria. Megasphaera elsdeni and Prevotella bryanti strains, yeast products containing Saccharomyces and Aspergillus LAB, may have beneficial effects on the rumen and intestinal tract.

Combining LAB and LUB may assist improve rumen conditions and promote dry matter intake, feed efficiency, and weight gain in cattle. Luebbe et al. (2013) demonstrated that the ADG and feed efficiency of cattle fed a DFM had numerical advantages. Ruminal fermentation activators generated from cultures of live microorganisms have recently attracted increased scientific and commercial attention (Galina, et.al.2010). It is possible to increase digestion and cell wall strength while also stabilizing ruminal microbes using an active ingredient rich in lactobacilli, yeasts, carbonated short-chain organic acids, and a low pH value (microbial addition). Elias and Herrera (2008) Increased proliferation of beneficial rumen microbes was a result of the addition of supplemental yeast. YC enhanced the quantity of anaerobic and cellulolytic bacteria in the rumen, which is a good thing. Yeast has been found to supply vitamins (particularly thiamin) that enhance the development of rumen fungus (Sun et.al.2021). As mutant Sc strains with impaired respiration could not increase the number of viable bacteria in the rumen, it seems that the capacity of various strains of Sc to remove oxygen from the rumen fluid is connected to their ability to promote bacterial counts (Li, et. al. 2016). Wheat straw + yeast had considerably greater total volatile fatty acid levels than wheat straw alone According to Xiao et.al. (2016), the ruminal pH and ammonia-N concentrations were not affected by SCFP treatment.



Fig 1. Probiotic Yeast: Effects on Ruminant Nutrition (Ghazanfar.et.al.2017)

Bacterial genus Saccharomyces cerevisiae

Kingdom	: Fungi
Phylum	: Ascomycota
Sub Phylum	: Saccharomycotina
Class	: Saccharomycetes
Order	: Saccharomycetales
Family	: Saccharomycetaceae
Genus	: Saccharomyces
Species	: Cerevisiae

The impact of Saccharomyces cerevisiae on the performance of ruminant animals

Feeding Routine

Feeding Routine However, dry matter had a negative impact on lamb development and rumen cellulolytic activity, but had no effect on feed intake. Increased feed intake has been linked to the use of probiotics(Moharrery and Asadi ,2009). The quantity of dry matter ingested by lambs and kids was unaffected by the addition of yeast culture, according to Titi. et al. (2008) (DM). According to Kumar et al. (2012), supplementing buffalo bull calves with 0.25 g/head/day of yeast culture had no effect on feed consumption. Awassi lambs fed diets enriched with SC consumed more DM and OM from barley straw on a daily basis, according to a study (Titi et al. 2008). According to Pienaar, et.al.(2012) When lambs were fed either a live yeast product or an ionophore, there was no significant difference in the amount of feed they consumed (P = 0.05). Male lambs were fed either steamflaked corn (CON), monensin (MO), or a baseline diet with live yeast (BYY) according to (Ding et al. 2008). (LY). The average daily consumption of monensin and live yeast supplements was not affected (ADI). While feeding at four doses (50, 100, 150, and 200 g/day) resulted in significant (P 0.01) increases in DM, ME, TN, and the response to Iraqi probiotics were higher with the greatest dosage of feeding, which is similar with the results of (Moharrery and Asadi ,2009) . According to Ding et al. (2008), the addition of dry yeast to goat kid meals has increased DMI, TDNI, and DCPI. When supplemented with 10 g Sc/cow/d, dairy cows receiving medium or high doses of nutritious concentrate had modestly higher DMI, according to (Ahlam ,2011). Only the DMI of dairy cows given a LYsupplemented feed differed from that of the control group.



Fig 1. The mechanism by which yeast increases production in ruminants is shown schematically [Xiao et.al. 2016; Wallace,1994]

Ratio of feed to live weight increase

More study is needed to better understand the function of yeast culture (Yc) and its mode of action in sheep and goats. Yalçın ,et.al(.2011) ; Ahmed and, Salah

10070

(2007) found that when goats' young were not given 2.5 or 5 grams of yeast per day, their average daily growth rate was much lower than when they were. Feed conversion ratios (FCR), average daily growth, live weight gain, and feed efficiency unaffected in lambs fed diets supplemented with 10% were Sc. Nathália, et.al. (2021) found that feeding Cows a yeast culture plus monensin increased weight growth in the animals. Lambs receiving a probiotic supplement gained more weight and had a lower feed conversion ratio (Hassan and Hassan,2009).

In calves given a full mixed diet with yeast culture, adding yeast culture at a rate of 20 kg/yeast/ton of feed had no influence on final weight, average daily growth, or feed conversion ratio. When feeding levels were raised (P 0.01), live weight growth was also enhanced (P 0.01), but the association between feeding levels and probiotics was not significant (Salim ,2009). An Iraqi probiotic or medicinal plant-supplemented lamb diet increased both live weight and feed conversion ratio significantly (Hassan and Hassan. 2010b; Mirzaei and Hari,2012) compared to an supplemented lamb diet. Iraqi probiotics, which included the bacterium Sc, increased body weight growth, feed conversion, and mortality in the study animals. To our knowledge, feeding lambs diets that included 0, 5, 15, 20, 30, and 40% percent yeast cultures had no effect on daily weight gain and feed conversion ratio (P = 0.10). This also holds true for feeding the lambs diets that had 0, 1, 5, 10, and 15% percentages of yeast cultures. Growth and feed conversion efficiency, as well as rumen fermentation indices, were improved when S. cerevisiae NCDC-49 was added (Özsoy ,et.al.2013).

Digestibility

Researchers have researched the effects of certain yeast cultures on ruminants' performance for over two decades. The effects of a supplemental yeast culture on rumen development have not been studied. Fermentation and rumen microbial populations may be affected by the rumen-viable facultative anaerobic yeast Saccharomyces cerevisiae (Sc). As a result of this study, it was shown that in-vivo DM digestibility, acid detergent fiber, nitrogen and in vitro DM digestibility did not differ significantly between the control and treatment groups. After 24 and 48 hours, Kholif and Khorshed (2006) found no difference in the digestion of (DM) when yeast was introduced to the diets of lactating buffaloes. This study by Abdel-Ghani (2004) found that goats fed the same meal, but supplemented with 3 or 6 grams of YC, absorbed more nutrients.

A similar trend emerged in feed consumption. Saccharomyces cerevisiae (Sc) has been demonstrated to help with ruminal digestion when taken as directed (Gaafar ,et.al.2009). Probiotics (yeast culture) had no effect on the digestibility of dry matter (DM), protein (CP), or NDF in Titi et al. (2008). Even additional study has shown that a mixture of OM and ADF may be more easily absorbed by lambs by adding yeast culture to the meal. According to Lascano et al. (2009) meat goats administered commercial probiotics metabolized more apparent dry matter, crude protein, neutral detergent fiber, and acid detergent fiber compared to the control group. Rumen digestion has been demonstrated to be impacted by yeast supplementation in studies (Han,et.al.2021). As reported by Lascano et al. (2009), adding yeast to a high-concentrate diet

(75.67 percent dry matter digestibility compared to 72.96 percent) improved the heifers' ability to digest dry matter. Between 73.65% and 73.97% of the population, For example, cellulose, hemicellulose and acid detergent fiber digestibility's increased when yeast supplementation was utilized by (Kholif and Khorshed ,2006).

Researchers observed that adding 15 g of baker's yeast per head per day to the 40:60 roughage feeding diet resulted in improved nutritional digestibility. Additionally, Kumar, et.al.(2012) discovered that the inclusion of 0.25 g/head/day of yeast culture in the buffalo bull calves' feed enhanced nutrient digestion. Ewes fed 60:40 concentrate: roughage diets supplemented with live dry yeast (5 or 7.5 g/h/d) had better DM, CP, CF, TDN, and DCP digestibility than lambs in the control group when Sc and (Sc + Ls) were included. Male lambs fed diets containing (0.5 and 1 kg/ton concentrate) Sc exhibited improved CF digestibility than the control group, whereas TDN and DCP did not differ between treatments. Probiotic supplementation with 20ppm and 0.1% probiotics increased DM, TDN and DP in lambs fed 18 to 22% protein, according to Sarwar and colleagues (2010) . Herawaty ,et.al.(2013) Beef cattle fed 0.5 percent Sc feed had higher nutritional digestibility than those fed a control diet, but poorer digestibility than those fed the supplement.

Characteristics of the Rumen Volatile Fatty Acids That Are Quickly Decomposable (VFA)

The rumen's metabolism and absorption of volatile fatty acids (VFAs) have a major chemical impact on rumen growth. Microbes naturally manufacture these (VFA) (Newbold, et.al.1995). Yeast supplementation had influence on ruminal (VFA) concentrations, according to Phesatcha, et.al.(2021) . When it comes to TVFA generation, Doleali and co-workers (2005) found that the use of ammonia was greater in the experimental groups than in the control groups (both at 8.40 and 8.40 mg/L, respectively). VFA concentrations in the rumen rose when Sc was added to a forage-enriched diet, according to Galip ,(2006). Between 24 and 48 hours, Kholif and Khorshed (2006) found no change in the concentrations of acetate, propionate, or butyrate (Lila,etal.2004) . TVFA content in rumen fluid increased from 91.26 to 103.34 mmol/l at 3 hours after feeding in Sc. compared to the control diet. Sc (Doležali, et.al.2005)] boosted TVF's concentration.

Longuski and colleagues ,2009 Sarwar, et.al.2010) g/cow/d) found no effect of yeast supplementation on total ruminal VFA or acetate concentrations (Stewart and Smith.2005; Longuski, et.al .2009). A concentration of 102.3 mml of VFA was shown to be unaffected by yeast supplementation or meal supplementation by researchers (Bal and Gokus 2013). Adding yeast had no influence on the acetate or propionate concentrations in the 50% or 70% concentrates. The concentrations of ruminal acetate and propionate, on the other hand, reduced and rose. At 2 and 4 months, TVFA concentrations in live YC-fed infants were considerably higher (P 0.01)(Özsoy ,et.al.2013).

The Rumen PH

For cellulolytic bacteria, which cannot thrive at pHs of 6.0 and lower, rumen pH is significant factor in rumen activity (Longuski ,et. al .2009). а Sulfur supplementation had no effect on ruminal pH Ahmed and, Salah (2007). When Sc was employed, pH values decreased significantly (P> 0.05) according to Gaafar and colleagues (2009) The greatest and lowest ruminal pH values were seen in the 2.5 g Sc and control groups (Chaucheyras ,et.al.2008). The ruminal pH may be elevated by feeding dairy animals probiotics or yeast culture, which has been demonstrated in other research to have health benefits for dairy cows and goats. Kholif and Khorshed (2006) found no difference in ruminal NH₃-N content when yeast was added. Sheep given yeast culture had a PH that was lower and fluctuated around the lower limit than the control group, according to Doleali et al. (2005) who found that adding Sc to the diet improved ruminal pH and decreased lactate levels in cattle. Yeast culture had little effect on rumen pH (Guedes and others .2008). Dairy cows on high-concentrate diets were shown to have lower ruminal pH levels(Yalcin, et.al.2011) than those fed a lower-concentrate diet.

Animal Ruminants Exposed to Ammonia

Microbe protein degradation and non-protein nitrogen use have been linked to rumen ammonia concentrations. Broderick and others (2008). Adding Sc to hay to a steer diet had no influence on NH₃-N concentrations, according to Dawson et al. (1990). As Newbold et al. (1995) reported, sheep fed Sc. boosted their NH_3 contents. There was an increase in ruminal NH3-N concentrations 4 hours after Sc administration (Biricik and Yavuz, 2001). S. cerevisiae supplementation was linked to a rise in the rumen-to-stomach ratio and an increase in amino acid absorption by small intestinal microbes [68]. Yeast had little effect on the small intestine's absorption of nitrogen fraction and amino acids, according to Putnam et al. (1997). According to Hong and Gallagher (1994) adding yeast had no influence on ruminal NH3-N concentration. At 1 g/day, Stewart and Smith (2005) found that adding Sc had no effect on the NH3-N concentration. When it came to commercial live yeast cultures, Sc lowered the concentration significantly. Supplementation with 4 g/day of live yeast culture showed no impact on NH3-N (Tripathi, et.al.2008; Saeed,2011; Hassan and Mohammed ,2016). According to Fatma et al. (2010), adding Sc at a concentration of 0.50 percent raised the NH3-N concentration.

Conclusions

Saccharomyces cerevisiae boosted total daily intake and improved the rumen environment in all trials with high dosages of concentrate supplementation.

References

- 1. Abdel-Ghani AA. Influence of diet supplementation with yeast (Saccharomyces cerevisiae) on performance of Zaraibi goats. Small Ruminant Research. 2004; 52:223-229.
- 2. Ahlam RA. Utilization of Saccharomyce cerevisiae supplementation for feeding goats in South Sinai. Egyptian Journal of Nutrition and Feeds. 2011; 14(2):169-181.

- 3. Ahmed BM, Salah MS. Effect of yeast cultures an additive to sheep feed on performance, digestibility, nitrogen balance and rumen fermentation. Journal of (*Saccharomyces cerevisiae*) culture on NDF digestibility and rumen fermentation of forage sorghum hay in Nubian goat's kids. Journal of Agriculture and Biological Science. 2007; 3:133-137.
- 4. Amin A. B., Mao, S. Influence of yeast on rumen fermentation, growth performance and quality of products in ruminants: A review Animal Nutrition .2021.Vol. 7, Issue 1, March, Pages 31-
- 5. Bal AM, Gokus S.Effects of live yeast supplementation on ruminal parameters and lactation performance of dairy cows fed medium or high levels of dietary concentrate.Kafkas Univercity of Veternal Fak Derg, 2013; 191:57-62.
- 6. Bayat A.R., Kairenius ,P. Stefański T., eskinen, H.L, Comtet-Marr,S. e Forano,E., Chaucheyras-Durand F.,
- 7. Biricik H, Yavuz HM. Effects of Saccharomyces cerevisiae yeast culture on milk production, milk composition Saccharomyces cerevisiae live cells on mixed ruminal microorganism fermentation in vitro. Journal of Animal Science. 2001; 82:1847-1854.
- 8. Broderick GA, Stevenson MJ,Patton RA.Effect of dietary protein concentration and degradability on response to rumen-protected methionine in lactating dairy cows Journal of Dairy Science. 2008; 92:2719-2728.
- 9. Dawson KA, Neuman KE, Boling JA. Effects of microbial supplements containing yeast and lactobacilli on roughage-fed ruminal microbial activities. Yeast Cultures in Ruminant Nutrition. Journal of Animal Science. 1990; 68(10):3392-3398.
- 10. Denev SA, Peeva TZ, Radulova P, Stancheva P, Staykova G, Beev G.(2007). Yeast cultures in ruminant nutrition. Bulgarian Journal of Agricultural Science.; 13:357-374.
- 11. Ding J, Zhou ZM, Ren LP, Meng QX. Effect of monensin and live yeast supplementation on growth performance, nutrient digestibility, carcass characteristics and ruminal fermentation parameters in lambs fed steam-flaked corn-based diets. Asian- Australian Journal of Animal Science. 2008; 21:547-554.
- Doležali P, Doležali J, Třináctŷ J. The effect of Saccharomyces cerevisiae on ruminal fermentation in dairy cows. Czechs.Journal of Animal Science. 2005; 50(11):503-510.Fuller R. A review: Probiotics in man and animals. Journal of Application Bacteriology1989.; 66:365-378.
- 13. Elías A, Herrera FR. Feed production through processes use of beneficial microorganism's evaluation active (MEBA), 2008, 8-13.
- 14. Fatma I, Emel G, Behic C, Mustafa SA, Ozcan BC, Esadsam P, *et al.* The effects of live yeast culture (Saccharomyces cerevisiae) on rum 71. en fermentation and nutrient degradability in yearling lambs Kafkas University of Veteriner Fafkas Dergisi. 2010; 16(5):799-804.
- 15. Gaafar HMA, Mohi AMA, Basiuoni MI, El-Riedy KF. Effect of concentrate to roughage ratio and baker's yeast supplementation during hot season on performance of lactating buffaloes. Slovak, Journal of Animal Science.2009;42(4):188-195.

- 16. Galina M, Pineda LJ, Morales R.(2010)Supplementation in dairy goats grazing with a slow release source of nitrogen (SLNN) with or without the addition of a probiotic lactic acid bacteria. III Congress of tropical animal production and symposium focal. Havana. Cuba,.
- 17. Galip N. Effect of supplemental yeast culture and sodiubicarbonate on ruminal fermentation and blood variables in rams. Journal of Animal Physiological and Animal Nutrition. 2006; 90:446-452.
- 18. García, G. A. R., & Rivas, Y. C. (2022). The circular economy and environmental training as care of the common house. International Journal of Life Sciences, 6(1), 1–9. https://doi.org/10.53730/ijls.v6n1.3148
- 19. Ghasemi E, Khorvash M, Nikkhah A. Effect of forage sources and Saccharomyces cerevisiae (Sc47) on ruminal fermentation parameters. African. Journal of Animal Science. 2012; 42(2):0375-1589.
- Ghazanfar, S., Nauman, K., ftikhar, A and Muhammad, I. Probiotic Yeast: Mode of Action and Its Effects on Ruminant Nutrition. Yeast - Industrial Applications.2017
- 21. Giler-Medina, P. X., & Alcívar-Castro, E. J. (2022). Variational thinking in the academic performance of High School students. International Journal of Health Sciences, 6(S5), 666–678. https://doi.org/10.53730/ijhs.v6nS5.8701
- 22. Guedes CM, Goncalves D, Rodrigues MAM, Dias A.Effects of a Saccharomyces cerevisiae yeast on ruminal fermentation and fiber degradation of maize silages in cows. Journal of Animal and Feed Science Technology. 2008; 145:27-40.
- 23. Han, G., Gao, X., Duan, J., Zhang, H., Zheng, Y., He, J., Huo, N., Pei, C., Li, H., Gu, S. Effects of yeasts on rumen bacterial flora, abnormal metabolites, and blood gas in sheep with induced subacute ruminal acidosis .Animal Feed Science and Technology Vol. 280,. 2021, 115042.
- 24. Hassan SA, Hassan KM. Effect of different levels of blood parameters of karadi lambs. Iraqi Journal of agriculture Science. 2009b.; 40(1):168-178.
- 25. Hassan SA, Hassan KM Effect of graded levels of rumen degradable nitrogen and Nigella Sativa on daily intake, live weight gain, feed conversion ratio and some blood parameters of karadi lambs. 7th Scientific Conference for Agriculture Research of Iraq. 2009a.; 14(5):194-204.
- 26. Hassan SA, Hassan KM. (2010b) Response of Karadi lambs to the Rosemary officinal supplementation fed with either alkali treated or untreated barley straw basal diets. Egypt. Journal of Application Science. ; 25:529-541
- 27. Hassan SA, Hassan KM. Effect of Nigella sativa and Rosemary officinalis supplementation on daily intake, live weight gain and carcass characteristics of Awassi lambs. Egyptian Journal of Nutrition and Feeds 2010a.;12(1):65-73.
- 28. Hassan SA, Hassan KM. The effect of supplementation of medicinal plants and probiotic on growth rate and some blood parameters of karadi lambs. Egyptian Journal of Nutrition and Feeds. 2009; 12(1):53-63.
- 29. Hassan SA, Mohammed SF. Effect of saccharomyces cerevisiae supplementation on rumen characteristics in Awassi lambs fed diets with different roughage to concentrate ratios. Iraqi Journal of Agricultural Science.2016; 47:1-11.
- 30. Hassan SA, Saeed AA. Effect of feeding different levels of dietary protein and addition of baker's yeast (Saccharomyces cerevisiae) on Awassi lambs

performance.3-Blood parameters. The second scientific conference for Agriculture College, University of Kufa. 2012; 18-19.

- 31. Herawaty RN, Jamarun M, Zain A, Ningrat RWS. Effect of supplementation Sacharonyces cerevisiae and leucaena leucocephala on low quality roughage feed in beef cattle diet. Pakistan Journal of Nutrition. 2013; 12(2):182-184.
- 32. Hong PH, Gallaghe JR. The effects of yeast supplement on digestibility of low quality roughage fed to sheep. Protocol Australia Soc. Animal Production. 1994; 20:398.
- 33. Hua-wei L, Jian-ming Z. Utilization of Chinese herbal feed additives in animal production Agricultural Sciences in China2011.; 10 (8):1262-1272.
- 34. Ismaiel AM, Ali EF, Abou-Ganema II. Effect of Tonilisat and Roemin W2 Supplementations on the Performance of Lambs. Journal of Biological Science.2010; 6:4 150.
- 35. Kamal R, Dutt T, Singh M, Nandan D, Patel M, Choudhary L. Effect of live Saccharomyces cerevisiae (NCDC-49) supplementation on growth performance and rumen fermentation pattern in local goat. Journal of Applied Animal Research. 2013; 3:41.
- 36. Khadem AA, Pahlavan M, Afzalzadeh A, Rezaeian M.Effects of live yeast Saccharomyces cerevisiae on fermentation parameters and microbial populations of rumen, total tract digestibility of diet nutrients and on the in situ degradability of alfalfa hay in Iranian Chall sheep. Pak. Journal of Biological Science. 2007; 10(4):590-597.
- 37. Khatta I. M, Abdel-Wahed, A.M., Khattab, I.M., Z, Uchenna Y. A. El-Keredy, A, Mabrouk, Z. Effect of dietary probiotics supplementation on intake and production performance of ewes fed Atriplex hay-based diet Livestock Science Vol 237, July 2020, 104065
- 38. Kholif SM, Khorshed MM. Effect of yeast or salinized yeast supplementation to rations on the productive performance of lactating buffaloes. Egyptian Journal Nutrition and Feeds, 2006; 9:193.
- 39. Krehbiel CR, Rust SR, Zhanga G, Gilliland SE. Bacterial direct-fed microbials in ruminant diets: Performance response and mode of action. Journal of Animal Science. 2003; 81:120-132.
- 40. Kumar DS, Rama PJ, Raghava RE, Sarja RK. Effect of yeast culture supplementation on nutrient utilization in Graded Murrah buffalo bull calves Livestock Research for Rural Development This paper formed part of the reproductive performance in ruminants. Fac. Of Agricuture Ain-Shams University. Cairo, Egypt. Journal of American Science. 2012; 8:2.
- 41. Lascano GJ, Zanton GI, Heinrichs AJ. Concentrate levels and Saccharomyces cerevisiae affect rumen fluid- associated bacteria numbers in dairy heifer's. Journal of Livestock Science. 2009; 126(1-3):189-194.
- 42. Li S., Yoon I., Scott M., Khafipour E., Plaizier J. Impact of Saccharomyces cerevisiae fermentation product and subacute ruminal acidosis on production, inflammation, and fermentation in the rumen and hindgut of dairy cows. Anim. Feed Sci. Technol. 2016;211:50–60. doi: 10.1016/j.anifeedsci.2015.10.01.
- 43. Lila ZA, Mohammed N, Yasui T, Kurokawa Y, Kanda S, Itabashi H. Effects of a twin of strain of Saccharomyces cerevisiae live cells on mixed ruminal microorganism's fermentation in vitro. Journal of Animal Science. 2004; 82(6):1847-1854.

- 44. Longuski RA, Ying Y, Allen MS. Yeast culture supplementation prevented milk fat depression by a short- term dietary challenge with fermentable starch. Journal of Dairy Science. 2009; 92:160-167.
- 45. Luebbe MK, Jenkins KH, Furman SA, Kreikemeier K. (2013)Effects of feeding microbial feed additives on growth performance and carcass traits of steers fed steam-flaked corn-based diets with wet distillers grains plus soluble, Nebraska Beef Cattle Report, , 66.
- 46. Maamour i , O and Ben Salem .M. Effect of yeast culture feed supply on growth, ruminal pH, and digestibility of fattening calves Food Sci Nutr. 2021 May; 9(5): 2762–2767.
- 47. Mirzaei F, Hari KR. Introduction of five well-known ayurvedic medicinal plants as feed additives on livestock's performance: A review. Global Journal of Research of Medical. Plants and Indigen medical .2012.;1(8):328-339.
- 48. Moharrery E, Asadi A. Effects of supplementing malate and yeast culture (Saccharomyces cerevisiae) on the rumen enzyme profile and growth performance of Lambs. Shahrekord University, Animal Science Agriculture, Shahrekord Iran. 2009; 115.
- 49. Mousa KM, El-Malky OM, Komonna OF, Rashwan SE. Effect of some yeast and minerals on the productive and reproductive performance in ruminants. Fac. Of Agricuture Ain-Shams University. Cairo, Egypt. Journal of American Science. 2012; 8:2.
- 50. Nathália T.S.Grigolett .o,Lucas G.Ghizzi,Larissa S.Gheller,Mauro S.da S. Dias,Alanne T.Nunes,Tássia B.P.Silva,Guilherme G.da Silva,Luiz F.Costa e Silva,Daniel N.Lobato,Francisco ó.Renn, P. Effects of a blend of live yeast and organic minerals or monensin on performance of dairy cows during the hot season Journal of Dairy Science2021, Vol, 104, Issue 11, Pages 11634-11645
- 51. Newbold CJ, Wallace RJ, Chen XB, McIntosh FM. Different strains of Saccharomyces cerevisiae differ in their effects on ruminal bacterial numbers in vitro and sheep Journal of Animal Science. 1995; 73:1811-1818.
- 52. Özsoy B, Yalcin S, Erodogan Z, Cantekin Z, Aksui T.Effects of dietary live yeast culture on fattening performance on some blood and rumen fluid parameters in goats. Revue Veterinary Medicine. 2013; 164(5):263-271.
- 53. Phesatcha, K., Burarat, P., Krittika, C. Metha W., Anusorn C. Changed Rumen Fermentation, Blood Parameters, and Microbial Population in Fattening Steers Receiving a High Concentrate Diet with Saccharomyces cerevisiae Improve Growth Performance. Vet Sci. 2021 Nov 28;8(12):294.
- 54. Pienaar GH, Einkamer OB, Vander H, Merwe J, Hugo A, Scholtz GDJ, Fair MD. The effects of an active live yeast product on the growth performance of finishing lambs. S. Afr.Journal of Animal Science, 2012, 42.
- 55. Putnam DE, Schwab CG, Socha MT, Whitehouse NL, Kierstead NA, Garthwaite BD. Effect of yeast culture in the diets of early lactation dairy cows on ruminal fermentation and passage of nitrogen fractions and amino acids to the small intestine. Journal Dairy Science. 1997;80:374-384.
- 56. Ruiz Garcia JR, Orozco HJN, Ibarra EP, Ortiz-Munoz G, Gaowth C, Olmedo J, Sanchez AEffect of an enhancer feed additive on lamb performance. Journal of Animal and Veterinary Advances 2011.; 10:332-333.
- 57. Saeed AA. Effect of level and degradability of ditary protein fed without baker, s yeast (saccharomyces cerevisiae) on Turkish a wassi lamb's

performance. College of Agriculture, Baghdad University. PhD. Thesis, 2011.

- 58. Salim HJH. . The effect different levels of feeding on karadi lambs response to local Iraqi probiotics.Sulaimani University. 2009 Msc. Thesis,.
- 59. Sarwar M, Nasir M, and Muhammad A, Shahzad M.Traditionalversus high input feeding system: Impact on nutrients intake, blood dynamics, hormone profil, weight gain and economics in growing lambs. Egyptian Journal of Sciences. 2010; 5(1):127-145.
- 60. Science . 2015.Vol ,98, Issue 5, May Pages 3166-318.
- 61. Seo JKK, Seon-Woo MH, Kim D, Santi DK, Kam K, Jong H. Direct-fed Microbials for ruminant animals. Asian-Aust. Journal of Animal. Science. 2010; 12:1657-1667.
- 62. Shing field K.J. Effect of camelina oil or live yeasts(Saccharomyces cerevisiae) on ruminal methane production, rumen fermentation, and milk fatty acid composition in lactating cows fed grass silage diets Journal of Dairy
- 63. Soren N, Tripathi MK, Bhatt RS, Karim SA. Effect of yeast supplementation on the growth performance of Malpura lambs. Tropical Animal Health and Production.2013; 45(2):547-549
- 64. Stewart GS, Smith CP. Urea nitrogen salvages mechanisms and their relevance to ruminants, non- ruminants and man. Journal of Nutrition Research. Revew. 2005; 18:49-62.
- 65. Sun, X, Yue W., Erdan W, Shu Z, Qianqian W., Yan Z., Yajing W., Zhijun C., Hongjian Y., Wei W, and Shengli Li.2021.Effects of Saccharomyces cerevisiae Culture on Ruminal Fermentation, Blood Metabolism, and Performance of High-Yield Dairy Cow Animals (Basel). Aug; 11(8): 2401
- 66. Suryasa, I. W., Rodríguez-Gámez, M., & Koldoris, T. (2021). Get vaccinated when it is your turn and follow the local guidelines. *International Journal of Health Sciences*, 5(3), x-xv. https://doi.org/10.53730/ijhs.v5n3.2938
- 67. Titi H, Dmour RH, Abdullah AY. Growth performance and carcass characteristics of Awassi lambs and Shami goat kid culture in their finishing diet. Journal of Animal Science.2008; 142:375-383.
- 68. Tripathi MK, Karim SA, Chaturvedi OH, Verma DL. Effect of different liquid cultures of live yeast strains on performance, ruminal fermentation and microbial protein synthesis in lambs. Journal of dairy Science and Nutrition. 2008; 92:631-639.
- 69. Wallace RJ. Ruminal microbiology, biotechnology and ruminant nutrition: progress and problems. Journal of Animal Science.1994; 72:2992-3003.
- 70. Xiao JX, Alugongo GM, Chung R, Dong SZ, Li L, Wu S, Yoon ZH, Cao JZ. Effects of Saccharomyces cerevisiae fermentation products on dairy calves: Ruminal fermentation, gastrointestinal morphology, and microbial community. Journal of Dairy Science. 2016; 99(7):5401-5412.
- 71. Yalçın SY, Yalçın S, Pınar C, ArifO, Cemalettin B, Önder E. The nutritive value of live yeast culture (Saccharomyces cerevisiae) and its effect on milk yield, milk composition and some blood parameters of dairy cows Asian-Australia. Journal of Animal Science. 2011; 24(10):1377-1385.