

**How to Cite:**

Mehta, M., Sharma, T., Miyajiwala, F., & Chandini, D. (2022). Intrathecal clonidine versus intrathecal midazolam with hyperbaric bupivacaine for post operative analgesia in abdominal hysterectomy: A randomised study. *International Journal of Health Sciences*, 6(S1), 3877–3886. <https://doi.org/10.53730/ijhs.v6nS1.5707>

# **Intrathecal clonidine versus intrathecal midazolam with hyperbaric bupivacaine for post operative analgesia in abdominal hysterectomy: A randomised study**

**Dr Malini Mehta**

HOD and Professor, Department of Emergency Medicine, Shrimati Bhikiben Kanjibhai Shah Medical Institute and Research Centre, Sumandeep Vidyapeeth (An institute Deemed to be university), Piparia, Vadodara, Gujarat, India

**Dr Tejash Sharma**

Associate Professor, Department of Anaesthesiology, Shrimati Bhikiben Kanjibhai Shah Medical Institute and Research Centre, Sumandeep Vidyapeeth (An institute Deemed to be university), Piparia, Vadodara, Gujarat, India

**Dr Fatema Miyajiwala**

3<sup>rd</sup> year post graduate resident doctor, Department of Anaesthesiology, Shrimati Bhikiben Kanjibhai Shah Medical Institute and Research Centre, Sumandeep Vidyapeeth (An institute Deemed to be university), Piparia, Vadodara, Gujarat, India

**Dr Chandini D**

Senior Resident, Department of Anaesthesiology, Shrimati Bhikiben Kanjibhai Shah Medical Institute and Research Centre, Sumandeep Vidyapeeth (An institute Deemed to be university), Piparia, Vadodara, Gujarat, India.

**Abstract**---Objectives: This randomised study was conducted to compare the effects of intrathecal clonidine and midazolam with hyperbaric bupivacaine for postoperative analgesia in patients undergoing abdominal hysterectomy. Materials and Methods: 60 patients of ages between 40 and 60 years of ASA grade I/II undergoing abdominal hysterectomy were randomly divided into two equal groups. Group C and group M received 0.5% hyperbaric bupivacaine 3ml with either clonidine 45µg with 0.2 ml saline or midazolam 2.5mg respectively intrathecally. Onset and duration of sensory blockade, haemodynamic changes, duration of post operative analgesia, number of rescue analgesics and side effects if any, were observed. Results: There was statistically significant difference in the onset and duration of sensory block ( $p < 0.001$ ) between the two groups. Duration of post

operative analgesia was significantly longer in midazolam group (373.33±24.22 minutes) than in clonidine group (328.5±21.78 minutes). Conclusion: Spinal anaesthesia with 2.5 mg midazolam as an adjuvant to 3ml hyperbaric bupivacaine provides longer duration of post operative analgesia compared to 45µg clonidine with 3ml hyperbaric bupivacaine with better haemodynamic stability without any adverse effects.

**Keywords**-- Clonidine, hyperbaric bupivacaine, intrathecal, midazolam, post operative analgesia

## Introduction

The main aim of postoperative pain management is to decrease a patient's pain to a level which can be tolerated by the patient, with minimal or no associated suffering or distress<sup>(1)</sup>. It is of utmost importance that the anaesthetic and analgesic technique should be such that it provides optimal conditions for surgery and also reduces postoperative discomfort, morbidity and mortality<sup>(2)</sup>.

Spinal anaesthesia is frequently used in sub umbilical surgeries like lower abdominal surgeries, lower extremity orthopedic and arthroscopic surgeries <sup>(3)</sup>. It is used commonly due to simplicity, minimum skill implementation, optimal operative condition, unhampered airway patency, lower risk of aspiration, less intraoperative blood loss, minimal biochemical and metabolic changes secondary to the stress of general anaesthesia, continued analgesia in the post-operative period and minimal post-operative morbidity <sup>(3,4,5)</sup>.

Intrathecal adjuvants like clonidine, neostigmine, midazolam, magnesium sulphate and opioids are increasingly used for prolongation of postoperative analgesia with local anaesthetic. They also intensify the subarachnoid block and offer hemodynamic stability<sup>(6)</sup>. Clinical studies have suggested that intrathecal clonidine prolonged sensory block and decreased anaesthetic requirements during surgery and increased postoperative analgesia without any clinically significant side effects.<sup>(4,7)</sup>

With the discovery of benzodiazepine receptors in spinal cord, midazolam is used intrathecally for prolongation of analgesia along with local anaesthetics. Several studies showed that intrathecal or epidural midazolam produced a dose dependent modulation of spinal noiceptive processing in animals and humans and was not associated with neurotoxicity, respiratory depression or sedation.<sup>(8,9,10)</sup>

In this study, the analgesic efficacy of intrathecal clonidine is compared with intrathecal midazolam as an adjuvant to hyperbaric bupivacaine in abdominal hysterectomy.

## Objectives

- To compare the onset of sensory blockade by both drugs.
- To compare the duration of sensory blockade by both drugs.
- To compare duration of analgesia.

- To compare the number of rescue analgesia in 24 hours.
- To compare side effects/complications if any in both drugs.

### Materials and Methods

After permission and clearance from the ethical committee (SVIEC/ON/MEDI/BNPG18/D19189), this randomised study was conducted at Dhiraj (tertiary care) hospital in Department of Anaesthesiology, Vadodara, Gujarat, India. 60 patients between ages 40 and 60 years of Grade I or II of American Society of Anaesthesiologists (ASA) classification, posted for elective abdominal hysterectomy between 2019 to 2021 were included in the study. All the patients who participated in the study were explained clearly about the purpose and nature of the study. Written informed consent was obtained. Sample size is calculated using following formula:

$$n = 2 \left( \frac{\sigma (z_{1-\alpha/(2\tau)} + z_{1-\beta})}{\mu_A - \mu_B} \right)^2$$

$$1 - \beta = \Phi(z - z_{1-\alpha/(2\tau)}) + \Phi(-z - z_{1-\alpha/(2\tau)}) \quad , \quad z = \frac{\mu_A - \mu_B}{\sigma \sqrt{\frac{2}{n}}}$$

### Table/Figure 1: Sample Size

where:

- n is sample size
- $\sigma$  is standard deviation
- $\Phi$  is the standard Normal distribution function
- $\Phi^{-1}$  is the standard Normal quantile function
- alpha is Type I error
- $\tau$  is the number of comparisons to be made

$\beta$  is Type II error, meaning  $1-\beta$  is power.


(According to the above formula the sample size is 56 patients, but to make the study convenient we took the sample size as 60 patients)

Inclusion criteria:

- Patient willing to participate in the study.
- ASA I, II patients.
- Age of patients: 40-60 years.
- No past history of allergic reaction, sensitivity or other form of reaction to local anaesthetics of the amide type.

Exclusion criteria:

- Refusal of patient.
- Patients allergic to any drugs.
- History of seizure disorder.
- Patients with renal, hepatic, cardiovascular and respiratory diseases, neurological disorders and neuropathies.



**Table/Figure 2: CONSORT flow diagram**

Using chit method, 60 patients were randomly distributed into two equal groups of 30 each.

Group C received Inj 3.0ml bupivacaine (0.5% hyperbaric) + 0.3ml clonidine (45µg) + 0.2ml normal saline intrathecally.

Group M received Inj 3.0ml bupivacaine (0.5% hyperbaric) + 0.5ml midazolam (2.5mg) intrathecally.

### **Preoperative preparation**

Pre – anaesthetic check-up was done one day prior to the surgery. Patients were evaluated for any systemic diseases and laboratory investigations were noted. All patients were kept nil by mouth for atleast 8 hours before the surgery. The procedure of spinal anaesthesia and visual analogue scale was explained to all the patients. On the day of surgery, intravenous (i.v.) line was secured with 20 gauge cannula and preloading with 10ml/kg ringer lactate was done. On arrival of patient in the operating room standard monitors were attached. Echocardiography (ECG), non-invasive systolic blood pressure (SBP) and diastolic blood pressure (DBP), heart rate (HR) and arterial oxygen saturation (SpO<sub>2</sub>) were monitored and recorded. Patients were pre-medicated with Inj. glycopyrrolate 0.2mg and Inj. ondansetron 4mg intravenously.

Patients were positioned in the sitting position. After painting with povidone-iodine and alcohol under aseptic and antiseptic precautions and draping, 23 gauge spinal needle was inserted in the midline at L<sub>3-4</sub> interspace. Drug was injected over 10-15 seconds after free flow of clear cerebrospinal fluid. The patient was placed in supine position immediately after injection.

Time of onset of sensory and motor block was recorded. HR, SBP, DBP, SpO<sub>2</sub> and sedation was monitored every 5 minutes up to 15 minutes and then every 15 minutes till the surgery ended. Onset of sensory block was noted as the time


taken to loss of pinprick sensation at L<sub>1</sub> after intrathecal injection. The level of sensory block was determined using pin prick test by checking at 2 minutes interval until two consecutive levels of sensory block was identical (i.e., fixation of the level).

Sedation was assessed by 'four point sedation scale' (Modified Wilson sedation scale) Table/Figure 3).

Score	Description
1	Spontaneous eye opening [awake and alert]
2	Drowsy, responsive to verbal stimuli
3	Drowsy, arousable to physical stimuli
4	Unresponsive

**Table/Figure 3: Four point sedation scale**

Side effects and complications were noted and treated. Bradycardia defined as fall in heart rate < 60/minute and it was treated with Inj. atropine sulphate 0.6mg i.v. Hypotension defined as the fall in SBP  $\geq$ 20% from the base level and was treated with Inj. mephentermine 6mg i.v. Respiratory depression was defined as respiratory rate  $\leq$ 10 breath/minute. After the patients were shifted to recovery room, HR, SBP, DBP, SpO<sub>2</sub>, sedation and complications, if any was monitored for 24 hours. Duration of sensory was noted. Patient's pain score was assessed by visual analogue scale (VAS) <sup>(11)</sup>- Figure/Table 4. Duration of analgesia was considered from the time of intrathecal injection to when VAS  $\geq$ 4. Inj diclofenac sodium 1.5mg/kg i.v. was given for analgesia. Number of rescue analgesic in 24 hours were recorded.



**Table/Figure 4: Visual Analogue scale**

### Statistical Analysis

The collected data were analysed by unpaired student – t test, chi square test and results obtained in the form of range, mean and standard deviation. The probability value 'p' of less than 0.05 considered statistically significant.

## Results

This study was conducted to compare the effects of intrathecal clonidine and midazolam as an adjuvant to bupivacaine to assess the postoperative analgesia in total abdominal hysterectomy. 60 patients posted for total abdominal hysterectomy between the ages of 40 years and 60 years were divided equally into two groups. The demographic variables such as distribution of age, weight and ASA grading was statistically insignificant between both the groups ( $p > 0.05$ ).

**Table/Figure 5: Onset and duration of sensory blockade.**

	Group C	Group M	p-Value	Significance
	Mean $\pm$ SD (in minutes)	Mean $\pm$ SD (in minutes)		
Onset of sensory block at L1	2.47 $\pm$ 0.90	1.63 $\pm$ 0.67	$p < 0.001$	Highly significant
Duration of sensory block	258 $\pm$ 17.45	281.17 $\pm$ 14.30	$p < 0.001$	Highly significant

**Table/Figure 6: Duration of analgesia**

	Duration of analgesia (in minutes)		p-Value	Significance
	Group C	Group M		
Mean $\pm$ SD	328.50 $\pm$ 21.78	373.33 $\pm$ 24.22	$p < 0.0001$	Highly significant

The mean duration of analgesia was statistically significantly longer in group M than group C ( $p < 0.0001$ ) as seen in Table/Figure 6

	No of rescue analgesics in 24 hours		p-Value	Significance
	Group C	Group M		
Mean $\pm$ SD	2.77 $\pm$ 0.58	1.87 $\pm$ 0.73	$p < 0.0001$	Highly significant

**Table/Figure 7: No of rescue analgesics**

The mean number of rescue analgesics given in 24 hours was more in Group C than in group M ( $p < 0.0001$ ) as seen in Table/Figure 7.

Intraoperative sedation score, respiratory rate, arterial oxygen saturation was within normal limits and comparable between the two groups ( $p > 0.05$ ) intraoperatively and post operatively.

Intraoperative Complications	Group C	Group M
Hypotension	33.3%	6.6%
Bradycardia	10%	nil

**Table/Figure 8: Intraoperative complications**

Intraoperatively, significant hypotension and bradycardia was noted in group C compared to group M (Table/Figure 8) Postoperatively, the difference in mean systolic blood pressure, diastolic blood pressure and mean heart rate was statistically not significant ( $p>0.05$ ) between the groups. No side effects were observed post operatively.

## Discussion

Clonidine is centrally acting partial alpha 2 adrenergic agonist. It inhibits voltage gated  $\text{Na}^+$  channels and when administered intrathecally it potentiates post-operative analgesia by hyperpolarizing A-delta and C fibre in the substantia gelatinosa of the spinal cord <sup>(17)</sup>.

Intrathecal midazolam is a benzodiazepine agonist which binds to GABA A receptor leading to enhancement of GABA activity. There is a high density of GABA A receptors in lamina II of the dorsal horn of human spinal cord, possibly explaining the pain modulation effect of midazolam. Benzodiazepines suppress afferent evoked excitation in the substantia gelatinosa and motor horn. Benzodiazepines agonists exert an inhibitory effect on spinal sensory and motor excitability <sup>(18)</sup>.

In our study, it was found that the difference in demographic variables was statistically insignificant between both the groups ( $p>0.05$ ) which was in accordance with Suchita A Joshi et al (2011)<sup>(13)</sup>, Anjali Bhure et al (2011)<sup>(8)</sup>, Pingley et al (2012)<sup>(14)</sup>, Piyush Kumar Sengar et al (2016)<sup>(15)</sup>, Dr. T.chandra Kumar et al (2018)<sup>(5)</sup>, Gandhi Gunjan P et al (2018)<sup>(16)</sup>.

**Onset of sensory block:** In our study, it was observed that the mean onset of sensory block at L1 was earlier in group M than in group C ( $p<0.05$ ). Our findings correlated with the findings of Suchita A Joshi et al (2011)<sup>(13)</sup>, Gandhi Gunjan P et al (2018)<sup>(16)</sup> (Depicted in Table/Figure 5).

**Duration of sensory block:** The mean duration of sensory block in group M was significantly longer as compared to group C ( $p<0.001$ ). Gandhi Gunjan P et al (2018)<sup>(16)</sup>, also reported longer duration of sensory block in midazolam group than clonidine group ( $p<0.001$ ) (Depicted in Table/Figure 5).

**Mean duration of analgesia:** In our study, the mean duration of post-operative analgesia was longer in group M as compared to group C which was highly statistically significant ( $p<0.001$ ). Mean post-operative analgesic consumption in 24 hours in our study was statistically significantly less in group receiving midazolam compared to group receiving clonidine ( $p<0.001$ ) which correlates with the studies of Suchita A Joshi et al (2011)<sup>(13)</sup>, Dr. T.chandra Kumar et al (2018)<sup>(5)</sup> and Gandhi Gunjan P et al (2018)<sup>(16)</sup> (Depicted in Table/Figure 6, 9)

**Mean number of rescue analgesia in 24 hours:** In our study, the mean number of rescue analgesics given in 24 hours was more in Group C than in group M ( $p<0.0001$ ) which correlated with studies of Suchita A. Joshi et al<sup>(13)</sup> and Gandhi Gunjan P et al<sup>(16)</sup> (Depicted in Table/Figure 7,9).

SR NO	AUTHOR AND TYPE OF STUDY	STUDY POPULATION	TYPE OF SURGERY	DOSES OF DRUGS USED	DURATION OF ANALGESIA (Mean± Standard deviation)	NO OF RESCUE ANALGESICS
1	Suchita A. Joshi et al <sup>(13)</sup> Randomized, double blind, prospective, parallel group clinical trial.	n=50 Group BC: 25 Group BM:25	Lower abdominal surgery	Clonidine Midazolam	Clonidine: 296.60 ± 52.77 min Midazolam :391.64 ± 132.98 min	Less in BM group
2	Dr. T.chandra Kumar et al <sup>(5)</sup> . Double blinded, randomized case control study	n=60 Group BC: 30 Group BM:30	Lower abdominal surgeries	Clonidine 30 µg Midazolam	Clonidine: 306.17 mins Midazolam: 486.17 mins	More in BM group.
3	Gandhi Gunjan P et al <sup>(16)</sup> . Observational study	n=60 Group BC: 30 Group BM: 30	Hernia surgeries	Clonidine 30 µg Midazolam 2mg	Clonidine: 252.5±21.1 mins Midazolam: 351.6±39.1 mins	Less in BM group
4	Anjali Bhure et al <sup>(8)</sup> Prospective, randomized comparative study.	n=120 Group A(control):30 Group B(clonidine):30 Group C(fentanyl): 30 Group D (midazolam):30	Elective caesarian section	Clonidine 75 µg Midazolam 2.5 mg Fentanyl 25 µg	Clonidine: 426.70 ± 151.83 mins Midazolam 270.54 ± 36.22	*
5	P INGLE Y et al <sup>(14)</sup> Prospective randomized study.	n=75 Group A(control): 25 Group B(clonidine) :25 Group C (midazolam): 25	Lower abdominal and lower limb surgeries	Clonidine 75 µg Midazolam 2.5 mg	Clonidine: 420.50 ± 130.80 mins Midazolam: 290.54 ± 46.22	*
6	Piyush Kumar Sengar et al <sup>(15)</sup> . Randomised study.	n=40 Group C: 20 Group M: 20	Lower limb surgeries	Clonidine 75 µg Midazolam 2.5 mg	Clonidine: 8.69 ± 0.7 hrs Midazolam: 4.28 ± 0.92 hrs	*
7	Our study Randomised study.	n=60 Group C: 30 Group M: 30	Abdominal hysterectomy	Clonidine 45 µg Midazolam 2.5 mg	Clonidine: 328.5±21.78 minutes Midazolam: 373.33±24.22 minutes	Less in BM group

\*- Data not available.

**Table/Figure 9: Comparison of various studies which used intrathecal clonidine and midazolam.**



### Study Limitation

We could not measure the plasma level of drug. Study was restricted to only female patients due to the nature of the study. Sample size is limited, so there is scope to study on large group.

### Conclusion

It is concluded that addition of 2.5 mg midazolam to 3ml hyperbaric bupivacaine intrathecally, resulted in faster onset of sensory block, longer duration of sensory block as well as prolonged duration of post operative analgesia compared to 45µg clonidine with 3ml hyperbaric bupivacaine without any side effects. Patients in clonidine group had hypotension and bradycardia but was not significant.

### References

1. Wylie and Churchill-Davidson's, *A Practice of Anesthesia* 2003 7th ed. Healy TJ, Knight PR, eds. London: Arnold Press:1217
2. Patil SS, Kudalkar AG, Tendolkar BA. Comparison of continuous epidural infusion of 0.125% ropivacaine with 1 µg/ml fentanyl versus 0.125% bupivacaine with 1 µg/ml fentanyl for postoperative analgesia in major abdominal surgery. *J Anaesthesiol Clin Pharmacol.* 2018;34(1):29-34.
3. Patil M, Padmawar S. Comparative study of intrathecal bupivacaine with bupivacaine plus clonidine in lower limb surgeries. *International Journal of Contemporary Medical Research* 2018;5(2):B45-B48.
4. Penchalaiah C. H, Kumar M. A Comparative Study of Clinical Effects of Intrathecal 0.5% Bupivacaine with Clonidine Versus 0.5% Bupivacaine in Patients Undergoing Major Gynecological Surgeries. *Journal of Evidence based Medicine and Healthcare*; 2015 July 27;2(30) : 4439-46.
5. Chandra Kumar T, Senthil Kumar S. Post operative analgesic requirement and adverse effects of intrathecally administered adjuvants midazolam and clonidine with bupivacaine. *Indian Journal of Applied Research* 2018;8(8):11-12.
6. Kathuria B, Luthra N, Gupta A, Grewal A, Sood D. Comparative Efficacy of Two Different Dosages of Intrathecal Magnesium Sulphate Supplementation in Subarachnoid Block. *Journal of Clinical and Diagnostic Research.* 2014 Jun;8(6)
7. Tarekegn F, Hailekiros A, Eshetie S. Comparison of the postoperative analgesic effects of intrathecal Clonidine and Fentanyl as adjuvant: meta analysis of randomized control trials. *Journal of Anesthesia & Critical Care: Open Access.* 2019 April;11(2):52-58.
8. Bhure A, Kalita N, Ingley P, Gadkari C.P. Comparative study of intrathecal hyperbaric Bupivacaine with Clonidine, Fentanyl and Midazolam for quality of anaesthesia and duration of post operative pain relief in patients undergoing elective caesarean section. *People's Journal of Scientific Research.* 2012 Jan;5(1)
9. Schoeffler P, Auroy P, Bazin JE, Ttvci J, Woda A. Subarachnoid midazolam: histologic study in rats and report of its effect on chronic pain in humans. *Reg Anesth* 1991; 16: 329-32

10. Naguib M, el Gammal M, Elhattab YS, Seraj M: Midazolam for caudal analgesia in children: comparison with caudal bupivacaine. *Canadian Journal of Anesthesia*. 1995; 42(9): 758-64.
11. Bharathi K, Chandapet S, Sidda P. Comparative study of intrathecal bupivacaine with buprenorphine and bupivacaine with tramadol. *MedPulse International Journal of Anesthesiology*. September 2018; 7(3): 125-32.
12. Filos KS, Goudas LC, Patroni O, Polyzou V. Hemodynamic and analgesic profile after intrathecal clonidine in humans. A dose-response study. *Anaesthesiology*; 1994 September;81(3):591-01
13. Joshi SA, Khadke VV, Subhedar RD, Patil AW, Motghare VM. Comparative evaluation of intrathecal midazolam and low dose clonidine: efficacy, safety and duration of analgesia. A randomized, double blind, prospective clinical trial. *Indian J Pharmacol*. 2012 May;44(3):357-61.
14. Ingley P, Kalita N, Gadkari C.P, Bhure A. Effects of addition of clonidine & midazolam to intrathecal bupivacaine in patient's under-going lower abdominal & lower limb surgeries. *J MGIMS*; 2012 September; 17(ii): 34 – 39
15. Kumar Sengar P, Priyadarshini S, Setu P. To study efficacy and safety of addition of Intrathecal clonidine and midazolam to hyperbaric Bupivacaine added as an adjuvant for post operative analgesia. *International Journal of Current Research*. 2016 September; 8(09): 38847-50.
16. Gandhi G, Mehta M. An observational study to compare intrathecal midazolam and clonidine for post operative analgesia in patients undergoing elective hernia surgeries. *Indian journal of Anesthesia and Analgesia*. 2018; 5(3): 372-76
17. Víctor M. Whizar-Lugo, Juan C. Flores-Carrillo and Susana Preciado-Ramírez. Intrathecal Clonidine as Spinal Anaesthesia Adjuvant — Is there a Magical Dose? *Topics in Spinal Anaesthesia*. 2014 September.
18. Clavier, N., Lombard, M. C., & Besson, J. M. (1992). Benzodiazepines and pain: effects of midazolam on the activities of nociceptive non-specific dorsal horn neurons in the rat spinal