

How to Cite:

Das, S. ., & Kulkarni, P. (2022). Updated review on overview of dry needling. *International Journal of Health Sciences*, 6(S1), 5127–5140. <https://doi.org/10.53730/ijhs.v6nS1.5991>

Updated review on overview of dry needling

Subhanjan Das

Department of Physiotherapy, Garden City University, Bangalore

Dr. Prema Kulkarni

Assistant Professor, Department of Lifesciences, Garden City University, Bangalore

Abstract--Dry needling is a therapeutic method that involves inserting a filiform needle into a target area. Osteoarthritis is a degenerative joint disease that affects the articular cartilage, causes discomfort, and changes the biomechanics. Ling is a component of conservative or comprehensive physiotherapeutic management. Increased blood flow, metabolic changes, spontaneous electrical activity, pain gating, and other mechanisms of action of dry needling may generate therapeutic advantages in osteoarthritis. The history of dry needling is extensively discussed. Needling can be defined as superficial or deep needling depending on the depth of the needle insertion, and it can also be classified as trigger point dry needling, fascial needling, scar tissue needling, and so on. The mechanism of Dry Needling is explained by a variety of ideas. The local effects of dry needling were also explored in this review, as well as the mechanism of analgesic effects. The needles can be utilized to provide therapeutic current to the tissues in a variety of ways. It can be done with PENS (Percutaneous Electrical Nerve Stimulation) or ETOIMS (Electro Transcutaneous Electrical Nerve Stimulation) (Electrical Twitch Obtaining Intramuscular Stimulation). This review effectively brought numerous features of dry needling to light while also discussing the current therapeutic state of dry needling.

Keywords--dry needling, osteoarthritis, pain, needling, pens.

Introduction

Dry needling is the procedure of insertion of filiform needle into the target tissue for therapeutic benefit. The potential benefits of dry needling include deactivation of trigger point, reduction of pain, promotion of healing, and alterations in the neuro-myo-fascial continuum. Although osteoarthritis is a degenerative joint disease that primarily affects the articular cartilage, pain and altered biomechanics in osteoarthritis also affects other contractile and non-contractile

soft tissues. Mild to moderate osteoarthritis is treated conservatively, which includes analgesics, exercises, lifestyle modifications, and physiotherapeutic interventions directed to articular and soft tissues (Dunning et al., 2014; Calvo-Lobo et al., 2018).

In recent times there has been an increase in the use of dry needling as the part of comprehensive physiotherapeutic or conservative management. There are several mechanisms of action of dry needling including increased blood flow, biochemical changes, spontaneous electrical activity, pain gate etc which may produce therapeutic benefits in osteoarthritis. Figure 1 shows a typical scene of dry needling (Butts et al., 2021).



Figure 1: Typical scene of dry needling

History of dry needling

The idea of dry needling emerged from the usage of injectable anesthetic for treatment of musculoskeletal pain. In 1930s, Kellgren suggested the concept of tenderness and illustrated the pattern of pain referral from specific muscles. In 1942, Janet Travell et al. (1942) used the term "trigger points" to describe tender points and brought the concept of myofascial trigger points. Again in 1941, Brav and Sigmond put forward that pain can be relieved by simple technique of needling without injecting any chemical substance. This was the first time a published paper brought dry needling to attention. However Brave and Sigmond referred Churchill's published paper dated back to 1821 which mentioned the injection treatment of sciatica and lower back pain. In 1979, Lewit suggested that the conventional drug injection therapy comprise of 2 independent parts: the role played by the injected drug and the effect of needling (Lewit K., 1979). Soon after this number of researches on the effect of needling saw an upsurge. This method was named "dry needling" to discriminate it from the customary drug injection, commonly referred as "wet needling". Since then Dry Needling has been extensively used in therapeutic purpose, specially tendon lesions and pain caused by joint replacement surgery (Mayoral O., et al, 2013). Figure 2 gives a summary of evolution of dry needling.




Figure 2: Evolution of dry needling

Types of dry needling

Dry Needling can be classified in various ways. Based on the depth of the needle insertion, needling can be termed as superficial and deep needling, based on the structure that is being needled it can be classified as trigger point dry needling, fascial needling, scar tissue needling etc. The classifications are almost always overlapping. Listed below are a few popular types of needling. In *Trigger-point Dry Needling*, MTrPs are localized hypersensitive spots in a palpable taut band of muscle. (Gattie E., et al, 2017). Trigger points can be diagnosed by a cluster of three diagnostic criteria: a taut band, a hypersensitive spot, and referred pain. (Fernandez C, Dommerholt J. International Consensus on Diagnostic Criteria and Clinical Considerations of Myofascial Trigger Points: A Delphi Study. *Pain Medicine* 2018; 19: 142–150). In MTrP DN needles are inserted directly into the trigger point to mechanically disrupt it. Often a small twitch is felt/seen during the procedure, known as local twitch response (LTR). While traditionally an LTR was an essential component of DN, evidences suggest that needling is as effective without LTR (Perreault et al., 2017). In *Superficial Dry Needling* the dry needles are inserted 5-10 mm deep into the skin without going into the target tissue. Peter Baldry was the proponent of Superficial Dry Needling (SDN) and he argues that because the pain of MTrP is primarily because of hypersensitivity of nociception, it can be modulated by A delta stimulation, which is possible through insertion of needle through a short distance. (Griswold, D., et al, 2019; Baldry P., 2002). Because SDN does not go deep into the tissues it is a much safer alternative (Griswold, D., et al, 2019). In *Deep Dry Needling* (DDN) a longer needle is used to reach the target tissue and it is often manipulated upon insertion (Ceccherelli F., et al, 2002; Fernández-Carnero, Josué., et al, 2017). DDN in a lot of ways are similar to MTrP DN. However, DDN can be employed in tissues other than muscles. There have been evidences of effectiveness of DDN in patients having problems in spine and the pain is unbearable and intense (Boluk and Huma., 2016). *Intramuscular Stimulation* was developed by Chan Gunn around 1970's. Instead of regarding pain to be signals of tissue injury, IMS views pain because of nerve pathology. Compression of nerve via soft tissue, which Gunn calls 'radiculopathy model' is the cornerstone of this needling style. (C Chan Gunn, 2002). Another type of dry needling started by Dr Yun Tao Ma, *Integrative Dry Needling*, this school of dry needling involves needling of the nerves and reflex points (Ma, 2011). Dry Needling for Hypertonicity and Spasticity: DNHS® has been proposed by Pablo Herrero who describes the use of dry needling to create therapeutic benefits in neural as well as mechanical aspects of the neuromuscular continuum. This type of needling is used specifically in hypertonicity and spasticity. Other less popular needling schools include spinal segmental sensitization, Fu's subcutaneous needling etc (Gallego & del Moral, 2007).

Mechanism of dry needling

There are many theories (Zylstra, E., & Maywhort, K.R. 2017) that explain the mechanism of Dry Needling. Some of the most accepted ones are:

- Enhanced blood flow- it is said that 'needling of the trapezius and Achilles tendon' accelerates flow of blood at the localized, needled site.

- Subsided banding- it has been perceived that after undergoing Dry Needling banding of the target tissue gets dwindled and occurs restitution of ‘sarcomere length and endomysium spacing’.
- Ebbed spontaneous electrical activity- stimulation of ‘local twitch response of a trigger point’ tapers electrical activity at that site which may show up as normalcy getting restored at neuromuscular junction.
- Biochemical changes- After Dry Needling therapy (of an Active TP) there is seen a local suppression of high concentrations of H⁺ ions, ‘neurotransmitters (bradykinin, 5HT, NE, CGRP, and substance P), cytokines, and chemokines (TNF- α , IL-1b, IL-6, and IL-8)’.
- Pain gate- the thought has also been proposed that action of needle insertion may bring about ‘sensory and proprioceptive stimuli’ which may impact the ‘gate control mechanisms’ of diminishing of pain.

The summary of the above explained theories are given below (Figure 3)




Figure 3: The summary of possible mechanisms of Dry Needling

Local effects of dry needling

Injury from needling produces rapid local vasodilatation and increased capillary permeability (Nakazato et al., 2021) which helps in healing and analgesia. Injury potentials are generated and can persist and provide stimulation for days until the miniature wound heals. This stimulation produces synthesis of prostaglandins which increase vascular permeability. Mast cell damage causes the release of histamine and heparin causing vasodilatation. Pain is relieved as a result of improved perfusion and relief of muscle spasm causes by local effects of needling and somatovisceral reflexes (Langevin et al., 2006).

The local effects also include increased local cortisol levels which have catabolic effect on connective tissue stimulating tissue remodeling and scar tissue breakdown. (Shah et al., 2005). The needle, once inserted, can be manipulated by rotation and pistoning. Needle manipulation causes pulling of collagen fibers towards the needle and initiates specific changes in fibroblasts. The fibroblasts respond by changing shape. The transduction of the mechanical signal into

fibroblasts can lead to a wide variety of cellular and extracellular events leading to neuromodulation and healing (Langevin et al., 2006).

Analgesic effects

The analgesic effect of dry needling involves multiple mechanisms including immune, hormonal, and nervous systems. The stimulation of A δ blocks the pain transmission of slow C fiber, this effect may persist for hours after needling due to the injury potential. Stimulation of the sensory afferent A δ also activates enkephalinergic, serotonergic, and noradrenergic inhibitory systems. A delta primary afferent receptors project to marginal cells (M) which carries pinprick sensations in the lateral spinothalamic tract. At the same time the A delta primary afferent receptors project to enkephalinergic stalked cells (ST) in the spinal cord, which inhibit the cells of substantia gelatinosa of Rolando (SGR) and block the transmission of C pain. This is known as segmental modulation. The serotonergic system involves midbrain and medulla. The fast pain carrying fibers of lateral spinothalamic tract give collaterals to peri aqueductal gray matter (PAG) of mid brain. PAG in turn activates Nucleus Raphe Magnus (NRM) of medulla which stimulates ST cells and inhibits SGR (Bowsheer D, 1998; Cheng, 2014).

The noradrenergic system involves medulla and pons. The nucleus paragigantocellularis (PGC) of medulla influences locus coeruleus (LC) of pons or similar noradrenergic brainstem structures which controls firing of SGR and modulates pain. Also active are the Diffused noxious inhibitory control where direct input of A delta generated information to Reticularis Dorsalis (R) in medulla inhibits pain by its influence on spinal segment. Together all these systems of pain modulation are known as Opioid Mediated Analgesia System (OMAS) (Kitahata L. M. (1993; Al-Chaer, 2009).




Figure 4: The pain pathway, C and A δ , and its conduction route are depicted, from receptor to thalamus via 2nd order neuron, finally reaching cortex through 3rd order neuron




Figure 5: The Pain Pathway and its regulation

Trigger point and dry needling

According to Travell and Simons (Vas J., et al, 2004; Scharf HP., et al, 2006), a trigger point (TP) can be defined as a ‘hyper-irritable spot in a taut band of skeletal muscle or the fascia which is under pain when compression happens to it and the peculiar pain thus produced, is called as ‘tenderness, motor dysfunction, and/or autonomic phenomena’. A trigger point is termed ‘active trigger point’ when there is spontaneous pain. In contrast to it, a ‘latent trigger point’ qualifies on all elements mentioned above without the production of peculiar pain

Electromyographically a trigger point is a source of spontaneous electrical activity (SEA) coming from a dysfunctional motor end plate which experience a thousand fold increase in the acetyl choline level (Gerwin, 1999). TPs generally are known to exist in an acidic environment. Inserting needles into the trigger point creates a small contraction, called Local Twitch Reflex (LTR). The LTR is hypothesized to change in local environment, creating an increased pain threshold and thus reducing pain (Shah, 2021).

The twitch also results in stimulation of neural fibers (C and Ad). This, in turn, affects spinal cord bringing further inhibition to nociception and heterosynaptic inhibition (Zylstra, E., & Maywhort, K.R. 2017). The mechanism is schematically presented below (Figure 6).




Figure 6: The summary of the mechanism of dry needling

Process of dry needling

The needling process involves insertion, manipulation and withdrawal of the needling. Most often the insertion is done using a guide tube, which is a hard plastic tube through which the needle is tapped. The use of guide tube minimizes pain. The needles come in various sizes, both in diameter and length. The choice of needle length depends on the depth of the target tissue, which means for superficial dry needling a smaller (usually 13 mm needle) is used whereas a deep seated structure (e.g. piriformis) will need a much larger needle. The angle of insertion also has important implications. The needle should be angled towards the target tissue and away from a vulnerable tissue. Also, needling of superficial structures (superficial fascia/scar in the skin) needs more acute angle (eg 15 degree angle) (Dommerholt J., 2011).

Needle manipulation

Once inserted the needle can be manipulated for therapeutic purposes. Although chinese acupuncture has a wide variety of manipulation techniques, however, in case of dry needling, the manipulations can be summarized into 4 broad categories. 4 types of needle manipulations are:

- Pistoning: Pistoning is described as rapid movement of the needle in and out of the MTrP (Cerezo-Téllez and Ester et al., 2016).
- Fanning/Fishing: A variation of pistoning, this technique uses the inclination angle of the needle along with up and down movement. Rossi advocates manipulating the needle for at least 30 seconds with 1hz frequency (Rossi and Ainsley et al., 2017). Fanning has been claimed to increase the chances of locating the loci of trigger point and eliciting twitch response (Rowley N.C, 2001).
- Needle rotation: Once the needle is inserted through the skin to the target tissue, the needle can be rotated clockwise and anticlockwise. It has been found that needle manipulation produces winding and gathering of collagen around the needle. Within minutes of needle rotation pulling of collagen toward the needle induces an active cellular response in connective tissue fibroblasts up to several centimeters away from the needle (Dunning J. et al., 2014).
- This transduction of the mechanical signal into fibroblasts can lead to a wide variety of cellular and extracellular events leading to neuromodulation and healing. (Simons *et al* 2002)
- Pecking/ Peppering: In case of chronic inflammatory condition especially in the junction of periosteum and ligament/tendon, to accelerate the healing response an acute inflammation is created by 'Pecking' that is, tapping the needle on the bony area multiple times after insertion. The resulting acute inflammation brings forth healing (Taşoğlu et al., 2017).

Fishing




Figure 7: A typical depiction of fishing (needle manipulation)

Dry needling as a medium of current delivery

The needles inserted to the body is often used as an electrode to deliver therapeutic current to the tissues. This is usually done in two ways, first is the use of a current similar to TENS, which is termed as PENS (Percutaneous Electrical Nerve Stimulation) whereas the other one is termed as ETOIMS (Electrical Twitch Obtaining Intramuscular Stimulation). PENS is largely for pain relief. They are a biphasic continuous waveform, with frequencies either between 2-5Hz or 80-100Hz and a pulse duration ranging from 250 to 500 microseconds (Park et al., 2020; Plaza-Manzano et al., 2020). All forms of PENS are usually applied in a biphasic continuous waveform, at a low (2-5Hz) or high (80-100Hz) frequency with pulse duration ranging from 250 to 500 microseconds. It can be argued that the use of PENS involves other pain modulation pathways like Dorsal column and analgesic neurotransmitters like GABA in addition to the pathways already active from dry needling (Plaza-Manzano et al., 2020; Sayson & Hargens, 2008).




Figure 8: The action of Electrical Twitch Obtaining Intramuscular Stimulation on Pain Pathway

The ETOIMS on the other hand uses a low frequency stimulator on the needle to stimulate the needles to obtain visible contraction. ETOIMS uses a monopolar electrode to stimulate deep deep motor endplates (Annaswamy et al., 2011). It has been used successfully in chronic myofascial pain (Chu, J., 2004; Chu et al., 2017). ETOIMS also can potentially favorably affect muscle activation and performance by providing more effective activation of mechanoreceptors. The tool used for TOIMS is often the standard EMG stimulator, however specifically made needle stimulator is also available. The image below shows two such stimulators (Chu, 2004; Rainey, 2013).



Figure 9: EMG stimulators

Conclusion

This review has presented the scope of Dry needling elaborately. The potential benefits have been discussed with evidence. The beneficial effects of dry needling on mild to moderate arthritis has been discussed. The review highlighted that in recent days, the usage of dry needling in clinical practice has risen significantly,

either as single management or as a part of comprehensive physiotherapeutic schedule. The author emphasised on the process of dry needling and main 4 types of the manipulations practiced. Dry needling is also used a medium of conducting electrode during PENS and ETOIMS. Several aspects of dry needling are discussed to bring updated information about dry needling and to support the increasing demand and need in practice of physiotherapy.

References

- Perreault, T., Dunning, J., & Butts, R. (2017). The local twitch response during trigger point dry needling: Is it necessary for successful outcomes? *Journal of Bodywork and Movement Therapies*, 21(4), 940–947. <https://doi.org/10.1016/j.jbmt.2017.03.008>
- Ma, Y.-T. (2011). *Biomedical acupuncture for sports and trauma rehabilitation : dry needling techniques*. Churchill Livingstone.
- C Chan Gunn. (2002). *The Gunn approach to the treatment of chronic pain : intramuscular stimulation for myofascial pain of radiculopathic origin*. Elsevier Science.
- Lewit K: The needle effect in the relief of myofascial pain. *Pain* 1979;6:83–90.
- Cagnie B., Dewitte V., Barbe T. *et al*. Physiologic Effects of Dry Needling. *Curr Pain Headache Rep* 17, 348 (2013). <https://doi.org/10.1007/s11916-013-0348-5>
- Mayoral O, Salvat I, Martin MT, et al. Efficacy of myofascial trigger point dry needling in the prevention of pain after total knee arthroplasty: a randomized, double-blinded, placebo-controlled trial. *Evid Based Complement Alternat Med* 2013;2013:694941.
- Dommerholt J, del Moral OM, Grobli C (2006). "Trigger point dry needling" (PDF). *Journal of Manual & Manipulative Therapy*. 14 (4): E70–E87. doi:10.1179/jmt.2006.14.4.70E
- Itoh K, Hirota S, Katsumi Y, et al. Trigger point acupuncture for treatment of knee osteoarthritis: a preliminary RCT for a pragmatic trial. *Acupunct Med* 2008;26:17–26.
- Ceballos-Laita, L., Jiménez-del-Barrio, S., Marín-Zurdo, J., Moreno-Calvo, A., Marín-Boné, J., Albarova-Corral, M. I., & Estébanez-de-Miguel, E. (2021). Effectiveness of Dry Needling Therapy on Pain, Hip Muscle Strength, and Physical Function in Patients With Hip Osteoarthritis: A Randomized Controlled Trial. *Archives of Physical Medicine and Rehabilitation*, 102(5), 959–966.
- D'Ambrosia RD. Epidemiology of osteoarthritis. *Orthopedics* 2005;28(2 suppl):s201–5.
- Dunning, J., Butts, R., Mourad, F., Young, I., Flannagan, S., & Perreault, T. (2014). Dry needling: a literature review with implications for clinical practice guidelines. *Physical Therapy Reviews*, 19(4), 252–265. <https://doi.org/10.1179/108331913x13844245102034>
- Butts, R., Dunning, J., & Serafino, C. (2021). Dry needling strategies for musculoskeletal conditions: Do the number of needles and needle retention time matter? A narrative literature review. *Journal of Bodywork and Movement Therapies*, 26, 353–363. <https://doi.org/10.1016/j.jbmt.2020.12.003>
- Calvo-Lobo, C., Pacheco-da-Costa, S., Martínez-Martínez, J., Rodríguez-Sanz, D., Cuesta-Álvaro, P., & López-López, D. (2018). Dry Needling on the Infraspinatus

- Latent and Active Myofascial Trigger Points in Older Adults With Nonspecific Shoulder Pain. *Journal of Geriatric Physical Therapy*, 41(1), 1–13. <https://doi.org/10.1519/jpt.0000000000000079>.
- Carmona L, Ballina J, Gabriel R, et al. The burden of musculoskeletal diseases in the general population of Spain: results from a national survey. *Ann Rheum Dis* 2001;60:1040–5.
- Gattie E, Cleland JA, Snodgrass S. The Effectiveness of Trigger Point Dry Needling for Musculoskeletal Conditions by Physical Therapists: A Systematic Review and Meta-analysis. *Journal of Orthopaedic & Sports Physical Therapy*. 2017;47(3):133-149.
- Griswold, D., M. Wilhelm, M. Donaldson, K. Learman, and J. Cleland. “The Effectiveness of Superficial versus Deep Dry Needling or Acupuncture for Reducing Pain and Disability in Individuals with Spine-Related Painful Conditions: A Systematic Review with Meta-Analysis.” *Journal of Manual & Manipulative Therapy* 27, no. 3 (May 27, 2019): 128–40.
- Baldry P. Superficial versus deep dry needling. *Acupuncture in Medicine*. 2002;20(2-3):78-81.
- Boluk, Huma. “Comparison of Two Different Dry-Needling Techniques in the Treatment of Myofascial Pain Syndrome.” *Ağrı – The Journal of The Turkish Society of Algology*, 2016.
- Ceccherelli F, Rigoni MT, Gagliardi G, Ruzzante L. Comparison of Superficial and Deep Acupuncture in the Treatment of Lumbar Myofascial Pain: A Double-Blind Randomized Controlled Study. *The Clinical Journal of Pain*. 2002;18(3):149-153.
- Fernández-Carnero, Josué, Laura Gilarranz-de-Frutos, Jose Vicente León-Hernández, Daniel Pecos-Martin, Isabel Alguacil-Diego, Tomás Gallego-Izquierdo, and Aitor Martín-Pintado-Zugasti. “Effectiveness of Different Deep Dry Needling Dosages in the Treatment of Patients with Cervical Myofascial Pain: A Pilot RCT.” *American Journal of Physical Medicine & Rehabilitation* 96, no. 10 (October 2017): 726–33.
- Dunning J, Butts R, Young I, et al. Periosteal Electrical Dry Needling as an Adjunct to Exercise and Manual Therapy for Knee Osteoarthritis. *The Clinical Journal of Pain*. 2018:1.
- Perreault, T, S O Flannagan, M T Grubb, and R Grubb. “Mechanisms and Dose Parameters of Electric Needle Stimulation: Clinical Considerations – Part I.” *Acupuncture in Physiotherapy*, 2018, 10.
- Butts R, Dunning J. Peripheral and Spinal Mechanisms of Pain and Dry Needling Mediated Analgesia: A Clinical Resource Guide for Health Care Professionals. *International Journal of Physical Medicine & Rehabilitation*. 2016;04(02).
- Longhurst JC, Tjen-A-Looi S. Acupuncture Regulation of Blood Pressure. *International Review of Neurobiology Neurobiology of Acupuncture*. 2013:257-271.
- Casanueva B, Rivas P, Rodero B, Quintial C, Llorca J, Gonzalez-Gay MA. Short-term improvement following dry needle stimulation of tender points in fibromyalgia. *Rheumatol Int*. 2013 Apr 23.
- APTA. Physical therapists & the performance of dry needling: an educational resource paper. Alexandria, VA, USA: APTA Department of Practice and APTA State Government Affairs; 2012. [Google Scholar]

- NCBPTE. North Carolina Board of Physical Therapy Examiners, June 14, 2012. Position statement: intramuscular manual therapy (dry needling) [Google Scholar]
- Maitland GD. Vertebral manipulation, 2nd eDry Needling. London: Butterworths; 1968. [Google Scholar]
- Kaltenborn FM. Mobilization of the spine, 1st eDry Needling. Oslo, Norway: Olaf Norlis Bokhandel; 1970. [Google Scholar]
- McKenzie R, May S. The cervical and thoracic spine: mechanical diagnosis and therapy, 2nd eDry Needling, Vol. 1. Waikanae, NZ: Spinal Publications New Zealand; 2006.
- Manheimer E, Linde K, Lao L, Bouter LM, Berman BM. Meta-analysis: acupuncture for osteoarthritis of the knee. *Ann Intern Med.* 2007;146(12):868–77. [PubMed] [Google Scholar]
- Mavrommatis CI, Argyra E, Vadalouka A, Vasilakos DG. Acupuncture as an adjunctive therapy to pharmacological treatment in patients with chronic pain due to osteoarthritis of the knee: a 3-armed, randomized, placebo-controlled trial. *Pain.* 2012;153(8):1720-6.
- Ebrahim AHM, Ahmed GM, Elsayed E, Sarhan R. Effect of electroacupuncture TENS, stretching exercises, and prefabricated insole in patients with plantar fasciitis. *Sci J Al-Azhar Med Univ.* 2007;28(3):1–10. [Google Scholar]
- Domingo, A., Mayoral, O., Monterde, S., & Santafé, M. M. (2013). Neuromuscular damage and repair after dry needling in mice. *Evidence-Based Complementary and Alternative Medicine: ECAM*, 2013, 260806. <https://doi.org/10.1155/2013/260806>.
- Zylstra, E., & Maywhort, K.R. (2017). *Orthopaedic Physical Therapy Secrets*. (Third Edition).
- Núñez-Cortés, R., Cruz-Montecinos, C., Vásquez-Rosel, Á., Paredes-Molina, O., & Cuesta-Vargas, A. (2017). Dry needling combined with physical therapy in patients with chronic postsurgical pain following total knee arthroplasty: a case series. *Journal of orthopaedic & sports physical therapy*, 47(3), 209-216.
- Wang, G., Gao, Q., Li, J., Tian, Y., & Hou, J. (2016). Impact of needle diameter on long-term dry needling treatment of chronic lumbar myofascial pain syndrome. *American journal of physical medicine & rehabilitation*, 95(7), 483.
- Bennell, K. L., Buchbinder, R., & Hinman, R. S. (2015). Physical therapies in the management of osteoarthritis: current state of the evidence. *Current opinion in rheumatology*, 27(3), 304-311.
- Sehgal,P, Kumar.B, Sharma.M, Salameh A.A, Kumar.S, Asha.P (2022), Role of IoT In Transformation Of Marketing: A Quantitative Study Of Opportunities and Challenges, *Webology*, Vol. 18, no.3, pp 1-11
- Kumar, S. (2022). A quest for sustainium (sustainability Premium): review of sustainable bonds. *Academy of Accounting and Financial Studies Journal*, Vol. 26, no.2, pp. 1-18
- Nisha Nandal,Dr. Naveen Nandal,Dr Aarushi. (2020). Women Social Entrepreneurs: A growing trend in Indian Economy. *International Journal of Advanced Science and Technology*, 29(4s), 2246 - 2253.
- Sánchez Romero, E. A., Fernández-Carnero, J., Calvo-Lobo, C., Ochoa Saez, V., Burgos Caballero, V., & Pecos-Martín, D. (2020). Is a combination of exercise and dry needling effective for knee OA?. *Pain Medicine*, 21(2), 349-363.
- Fernández-De-Las-Peñas, C., Plaza-Manzano, G., Sanchez-Infante, J., Gómez-Chiguano, G. F., Cleland, J. A., Arias-Burúa, J. L., ... & Navarro-Santana, M. J.

- (2021). Is Dry Needling Effective When Combined with Other Therapies for Myofascial Trigger Points Associated with Neck Pain Symptoms? A Systematic Review and Meta-Analysis. *Pain Research and Management*, 2021.
- Rahou-El-Bachiri, Y., Navarro-Santana, M. J., Gómez-Chiguano, G. F., Cleland, J. A., López-de-Uralde-Villanueva, I., Fernández-de-Las-Peñas, C., ... & Plaza-Manzano, G. (2020). Effects of trigger point dry needling for the management of knee pain syndromes: a systematic review and meta-analysis. *Journal of Clinical Medicine*, 9(7), 2044.