

How to Cite:

Eyong Edu, M., Okon Asuquo, O., Ekpo Edet, E., & Abeshi Sylvester, E. (2022). Implanon uptake at the university of Calabar teaching hospital Calabar, Nigeria: A five year review. *International Journal of Health Sciences*, 6(S2), 7236–7243.
<https://doi.org/10.53730/ijhs.v6nS2.6752>

Implanon uptake at the university of Calabar teaching hospital Calabar, Nigeria: A five year review

Eyong Edu M

Department of Obstetrics and Gynaecology, University of Calabar, Nigeria

Okon Asuquo O

Department of Obstetrics and Gynaecology, University of Calabar, Nigeria

Ekpo Edet E

Department of Obstetrics and Gynaecology, University of Calabar, Nigeria

Abeshi Sylvester E

Department of Obstetrics and Gynaecology, University of Calabar, Nigeria
Corresponding author email: abeshisly2007@gmail.com

Abstract---Background: Implanon is a modern hormonal contraceptive with a long-acting and reversible effect. It is effective for up to 3 years and can be given during breastfeeding, thus preventing early pregnancy in about 99% of breastfeeding mothers. The current prevalence rate for contraceptive use in Nigeria is approximately 11%–15%. This rate is very low in spite of the high rate of sexual activity and widespread awareness of the various contraceptive methods among Nigerian adolescents and youths. Objective: To determine the socio-demographic profile of implanon acceptors, the side effects profile and reasons for discontinuation of the method in Calabar. Method: This was a five-year retrospective review of records of clients who accepted Implanon at the University of Calabar Teaching Hospital (UCTH), Calabar, Nigeria. Result: During the period under review, 311 clients accepted Implanon as a method of modern contraceptive in this tertiary facility. The mean age of users was $31.0 \pm 5.5SD$. Of this number, 97.7% were Christians while 2.3% were Muslims. 93.6% of the clients had at least secondary education and above, 52.7% of the clients had three or more children at the point of accepting the method. However, 80.0% of them desired future fertility. The commonest reported side effect was menstrual irregularities while the commonest reason for removal was the desire to conceive. Pear index was 0.064. Conclusion: Implanon is a highly effective and acceptable method of contraception though it commonly causes menstrual

irregularities and other side effects necessitating many users to discontinue usage. It is breastfeeding friendly, since most women usually resume sexual intercourse early in the puerperium and still desire to continue breastfeeding effectively. Education plays a major role in decision making about choice or need for contraception.

Keywords---Implanon, socio-demographic profile, menstrual irregularities, Calabar.

Introduction

Contraception plays a critical role in controlling unwanted pregnancies, fertility rate, population growth rate and maternal mortality rate which have been shown to be associated with each other.¹ Nigeria has a low contraceptive prevalence rate (15%) and a high (16%) unmet need for contraception.^{2, 3} She has a high total fertility rate (5.5%) and a high population growth rate (2.6%).^{4, 5} Nigeria is rated the fourth (4th) highest maternal mortality country in the world, with an estimated 814 deaths per 100,000 live births.^{4, 6}

Contraception interacts with the other factors by preventing women from having unplanned pregnancies and subsequently dying from pregnancy-related causes. It has also been known to improve the educational and economic status; especially that of girls and women.^{6, 7} There is a wide range of contraceptive methods available which include the traditional or folk methods, barrier methods, Intrauterine Contraceptive Devices (IUCDs), hormonal methods, emergency contraception and sterilization. The choice of contraceptive is mostly influenced by the education, poverty, awareness, ignorance, culture, religion, myth about the method, availability, cost and side effects of the type of contraception.^{8, 9}

Implanon is a modern hormonal contraceptive method, which is produced by organon and was first marketed in 1998.¹⁰ It is a long – acting reversible contraceptive (LARC) method. It is one of the contraceptive implants and is known to be safe, widely acceptable and highly effective.^{10, 11} Implanon is a progestin-only contraceptive, and consists of a non-biodegradable, single, sterile rod implant for subdermal insertion. It is 40mm in length and 2mm in diameter and contains an Ethinyl Vinyl Acetate (EVA) copolymer with 68mg of etonorgestrel surrounded by EVA copolymer skin.^{9, 12} It releases 60 - 70µg of etonorgestrel per day in the early (15th – 16th) weeks of insertion. This decreases to 35 - 45µg of etonorgestrel per day by the end of the first year of insertion. This further decreases to 30 – 40µg of etonorgestrel per day by the end of the third year of insertion.¹² Implanon can be removed at any time of the menstrual cycle but must be removed at the end of three (3) years after insertion. The onset of contraceptive efficacy is within 24 hours after insertion, if it was inserted within the first five days of the menstrual cycle. When it is inserted at any other time of the menstrual cycle, an alternative barrier method of contraceptive must be used for seven days before contraceptive efficacy is assured.¹³

The mechanisms of action of implanon are by suppression of ovulation, cervical mucus thickening and induction of endometrial atrophy.¹⁴ The advantages of

implanon include the fact that it is user independent and requires minimum medical follow-up after insertion.^{12, 13, 14} The side effects of implanon include menstrual irregularities (menorrhagia, metrorrhagia, reduced or scanty menstrual flow, shortened menstrual cycle, and amenorrhoea), weight gain, headache, raised blood pressure.^{12, 13, 14}

Implanon was introduced into the family planning unit of the University of Calabar Teaching Hospital (UCTH) in 2005. It was introduced at the same time with Jadelle, which is another contraceptive implant which is licensed to be used for five (5) years. They were both preceded by Norplant, another contraceptive implant which was withdrawn in same year 2005 shortly before these two implants were introduced. Implanon is usually inserted sub dermally in the inner aspect of preferably the left arm. The procedure does not require highly skilled medical personnel, as well trained nurses in family planning can carry out the procedure.

The objective of this study is to determine the sociodemographic profile of implanon acceptors in University of Calabar Teaching Hospital (UCTH) as well as the side-effects and reasons for discontinuation of use of the product.

Methodology

This was a five-year retrospective study of 311 clients who accepted implanon at the family planning clinic of the University of Calabar Teaching Hospital (UCTH), Calabar, Nigeria from January 1st, 2016 to December 31st, 2020. Data retrieved from their case records included socio-demographic profile, reproductive history, desire for future fertility, side effects profile and reasons for discontinuation of the method. Descriptive and analytical statistics was carried out on all data using SSPS statistics program. The result was presented in Tables as frequencies and percentages

Results

Table 1
Socio-demographic profile of Implanon acceptor

Age group (year)	Implanon N (%)
10-19	5 (1.6)
20-29	124 (39.9)
30-39	164 (52.7)
40-49	18 (5.8)
Religion	n (%)
Christianity	304 (97.7)
Islam	7 (2.3)
Educational status	n (%)
None	1 (0.3)
Primary	19 (6.1)
Secondary	61 (19.6)

Tertiary	230 (74.0)
Total	311 (100.0)

Table 2
Mode of contraceptive usage by Implanon acceptor

Mode of usage contraceptive	Implanon n (%)
Permanent	57 (18.3)
Temporary	249 (80.1)
Uncertain	5 (1.6)
Total	311 (100.0)

Table 3
Reproductive history of Implanon acceptors

Parity	Implanon n (%)
0A	24 (7.7)
2A	123 (39.5)
3 - 4A	146 (46.9)
_ > 5A	18 (5.9)
Total	311 (100.0)

Table 4
Prevention methods used by respondents for Implanon acceptor

Prevention method	Implanon n (%)
Nil	121 (38.9)
Condom	46 (14.6)
Jadelle	2 (0.6)
Implanon	10 (3.2)
Injectables	27 (8.7)
IUCD	18 (5.8)
Pills	17 (5.5)
Withdrawal	5 (1.6)
Natural	9 (2.9)
Total	311 (100.0)

Table 5
Complications/Side effects among Implanon acceptors

Complication/ side effect	Implanon N (%)
Amenorrhoea	41 (28.3)
Menstrual irregularities	86 (59.3)
Excessive weight gain	3 (2.0)
Localized device problems	2 (1.4)
Others	13 (9.0)
	69 (22.2)

No side effects	
No follow-up visit	97 (31.2)
Total	311 (100.0)

Table 6
Reasons for Discontinuation by implanon acceptors

Discontinuation Reasons	Implanon n (%)
Desire for pregnancy	30 (61.2)
Menstrual irregularities	13 (26.5)
Expiration	6 (12.3)
Total	49 (100.0)

Discussion

In this study, 311 women accepted implanon as their family planning method of choice. Only 5 (1.6%) of the clients were teenagers although implanon has been found to be suitable for teenagers. There is no record of teenagers' uptake of implanon in similar studies in the country except that done in Port Harcourt¹⁵, Ilorin¹⁶. This is contrary to what obtains in many European and American countries where adolescents form a significant proportion of users of contraception in general and implants in particular,^{17, 18, 19} The low uptake of adolescent contraceptive in Nigeria is not unconnected with the fact that sex education is hardly taught in our schools and the mention of sex even at home by parents or siblings is like a taboo.²³ There are no hospital or school based counselling units in Nigeria to cater for the needs of the teenagers and /or adolescents. Even where such facilities exist, there will be low patronage due the stigmatization. It is generally believed in Nigeria that the teenagers and adolescents have no business engaging or discussing sexual issues. It is considered a taboo in most communities.²³

Table I shows the mean age of Implanon users was 31.0 ± 5.5 SD with a range of 19 – 49 years. Majority (52.7%) of the implanon users belonged to the 30 – 39 years' age group. Table 11 shows the proportion of implanon users who choose the method as a permanent form of contraception 57(18.3%); 249(80.1%) choose it as a temporary form of contraception while 5(1.6%) were uncertain about their desire for future fertility.

In our study, 97.7% were Christians and 2.3% were Muslims. In a similar study carried out in Ilorin, Nigeria 52.3% were Christians and 46.6% were Muslim. This could be explained by the fact that Calabar is located at the southern part of Nigeria, where there are predominantly Christians. On the other hand, Ilorin is located in the North- central part of Nigeria, where there are high proportion of both Christians and Muslims from the Yoruba tribe, of almost equal proportion.

Majority of the clients, 230 (74%) had received tertiary education, 61(19.6%) had secondary education, 19(6.1%) had primary education while 1(0.3%) had no formal education. Among the implanon clients, 93.6% had at least secondary level of education and above. This is similar to studies done in Port Harcourt, Nigeria (95.8%) and Ilorin (88.6%) whereas those with at least secondary level of

education constituted majority of the implanon acceptors. A higher level of education has been found to be strongly associated with higher use of modern family planning methods.^{20, 21, 22, 23}

Table III Shows the number of living children the clients had prior to insertion of the implant. 24(7.7%) had no living child, 123 (39.5%) had one or two, 146(46.9%) had three or four, 18(5.8%) had five or more living children. During the study period, 145(46.6%) of the clients reported side effects. The commonest reported side effect was menstrual irregularities which occurred in 86(61.4%) clients. This was also the commonest side effect reported in Port Harcourt¹⁵, Ilorin¹⁶ and Benin City.⁶ It was noted that 97(31.2%) of the clients did not report for follow – up during the period of the study. Thus, it cannot be confirmed whether they had side effects or not. This was a major constraint of this study, as almost one-third of the acceptors were lost to follow up.

Table IV shows previous methods of contraception used by implanon acceptors. 121(38.9%) had never previously used any method of contraception. The previous methods of contraception used and their distribution are clearly shown in this table. During the period under review 49(15.8%) clients discontinued implanon. The commonest reason for discontinuation was desire for pregnancy in 30(61.2%) followed by menstrual irregularities in 13(26.5%). Table V shows that 145 of the implanon acceptors reported side effects during the study period. Among the commonest reported side effect was menstrual irregularities (59.3%) followed by amenorrhoea (28.3%). Weight gain was reported by 2.0% and localized implant problems reported by 1.4%. Other reported side effects included headache, dizziness, increased blood pressure, skin changes, abdominal bloating. Only 69(22.2%) of the clients had no side effect. However, 97(31.2%) of the clients did not report for any follow-up visit.

It was noticed during this study period that whereas 86(27.7%) reported having menstrual irregularities, only 13(4.2%) discontinued implanon due to this complaint. Menstrual irregularities are well-known complications of all forms of progestogen-only contraceptives^{21,22}. However, with adequate pre- and post-insertion counselling, it is found to be more tolerable. In addition, it is found to remit within six months of onset of contraception or with treatment. ^{20, 23} Majority of the implanon users (61.1%) had previously used another method of family planning. The commonest previous method of family planning used was Implanon (18%) followed by condom (5.1%).

Table VI shows that 49(15.8%) of the clients discontinued Implanon during the period under review. Out of these, 30(61.2%) discontinued the method due to the desire to conceive, 13(26.8%) discontinued due to menstrual irregularities and 6(12.2%) discontinued due to expiration of the method. It was also noted during the study period that although 52.7% of the clients had three or more children, 80.1% of them had desire for future fertility. This agrees with previous study done in Benin City²⁰ and Ibadan²². It is noticed that Implanon is much more commonly used as a temporary method of contraception than a permanent method.

Conclusion

Implanon is a safe, highly effective and reversible hormonal contraceptive implant. The commonest side effect is menstrual irregularities but the commonest reason for discontinuation is desire for pregnancy in the University of Calabar Teaching Hospital (UCTH), Calabar. During the period under review, only one woman was confirmed pregnant and this was confirmed after 1 year 10 months of use. It can be confirmed to be true failure of the contraceptive method since pregnancy occurred remote from insertion and was confirmed by a blood pregnancy test. The Pearl index was 0.064 which shows that it is a highly effective contraceptive method. Adequate pre - and post - insertion counseling would help to increase its uptake, reduce its discontinuation and eventually reduce maternal mortality ratio.

References

1. Etuk S. J., Ekanem A. D. Knowledge, attitude and practice of family planning amongst women with unplanned pregnancy in Calabar, Nigeria. *Nigerian Journal of Physiological sciences*. 2003; 18: 65 – 71
2. National Population Commission (NPC) and ICF macro (2009) Nigeria Demographic Health Survey 2008: Key findings. Calverton, Maryland, U.S.A: NPC & ICF macro. 11: 63 – 86
3. Austin A. Unmet Contraceptive need among married Nigerian women: an examination of trends and drivers. *J contraception* 2015 Jan; 91 (1) : 31 – 38
4. Canning D., Schultz T. P. The economic consequences of reproductive health and family planning. *The Lancet*. 2012;380: 165 – 171
5. Schultz T. P., Shareen J. Family planning and women's and children's health: consequences of an outreach program in Matlab, Bangladesh. *Demography*. 2013; 50 (1): 149 – 180
6. Okonofua F. and Odunsi K. (eds) *Contemporary Obstetrics and Gynaecology for Developing Countries*; 2003. WHARC, Benin City, Nigeria. 110 – 119.
7. Association of Reproductive Health Professionals (2008). *History of subdermal contraceptive implants*. Association of Reproductive Health Bulletin.
8. Monjok E, Smesny A, Ekabua JE. Contraceptive practices in Nigeria: Literature review and recommendation for future policy decisions. *Open access contraception* 2010; 1: 9 – 22
9. Power J, French R, Cowan FM. Subdermal implantable contraceptives versus other forms of reversible contraceptives or other implants as effective methods of preventing pregnancy. *Cochrane Database of Systematic Reviews*. 2007: 3.
10. Adekunle AO. Recent advances on Contraceptive development. In: Ojule JD, Oranu EO, Enyindah CE. *Experience with implanon in Southern Nigeria*. *Journal of Med and Med Sciences*, 2012; 3(1):710 – 14.
11. Balogun OR, Olaomo N, Adeniran AS, Fawole AA. Implanon subdermal implant: an emerging method of contraception in Ilorin, Nigeria. *Journal of Med and Biomedical Sciences*, 2014; 3(1):1 – 5
12. Mestad R, Secura G, Allsworth JE, Madden T, Zhau Q, Peipert JF. Acceptance of long-acting reversible contraceptive methods by adolescent participants in the contraceptive CHOICE project – *Contraception* 2011; 84:493 – 8.

12. Guazzelli CA, de Queiroz FT, Barbieri M, Torloni MR, de Araujo FF. Etonorgestrel implants in adolescents: Evaluation of clinical aspects. *Contraception* 2011; 83:336 – 9
13. Lewis LN, Doherty DA, Hickey M, Skinner SR. Implanon as a contraceptive choice for teenage mothers: A comparison of contraceptive choices, acceptability and repeat pregnancy. *Contraception* 2010; 81:421 – 6.
14. Marrone G, Abdul-Rahman L, De Coninck Z, Johansson A. Predictors of Contraceptive use among female adolescents in Ghana. *Afr J Reprod Health* 2014; 18:102 – 9.
15. Ankomah A, Anyanti J, Oladosu M. Myths, misinformation and communication about family planning and contraceptive use in Nigeria. *Open Assess J Contracept* 2011; 2: 95 – 105
16. Palamuleni ME. Socio-economic and demographic factors affecting contraceptive use in Malawi. *Afr J Reprod Health* 2013; 17:91 – 104
17. Aisien AO, Enosolease ME. Safety, efficacy and side-effects of implanon, a single rod implantable contraceptive (etonorgestrel) in University of Benin Teaching Hospital. *Niger J Clin Pract.* 2010; 13(3):331 – 5.
18. Akadri AA, Odelola OI. Progestogen – only injectible contraceptive: Acceptor prevalence and client experience at Sagamu, Nigeria. *Niger Postgrad Med J* 2017; 24:178 – 81
19. Adeyemi AS, Adekanle DA, Progestogen – only injectable contraceptive: Experience of women in Osogbo, Southwestern, Nigeria. *AM Afr Med* 2012; 11: 27 – 31.
20. Hou MM, McNicholas C, Creinin MD. Combined oral contraceptive treatment for bleeding complaints with etonorgestrel contraceptive implant: A randomized control trial. *Eur J Contracept Reprod Healthcare* 2016; 21:361 – 66.
21. Friedlander E. Therapeutic options to unscheduled vaginal bleeding associated with long-acting reversible contraception. *Obstetrics and Gynaecology Clinics of North America* 2015; 42: 593 – 603.
22. Adebola O, Imran O, Morhason – Bello, Michael A, Okuniola, Adeyemi O, Adekunle. Profile of Implanon acceptors and pattern of side effects. *J Reprod Med* 2015, Mar; 26(1): 46 – 52
23. Abeshi, Sylvester E; Eyong, Edu M; Ekenjoku, A J; Ekpo, Edet E -Current trend in contraceptive demand at the university of Calabar teaching hospital, Calabar, Nigeria. *Asian journal of science and technology* 11 (05); 10972-10974, 2020