

How to Cite:

Tiwari, S. P. (2022). Application of nanomedicine in the treatment of melanoma: Advances and prospects. *International Journal of Health Sciences*, 6(S1), 12124–12134.
<https://doi.org/10.53730/ijhs.v6nS1.8038>

Application of nanomedicine in the treatment of melanoma: Advances and prospects

Sandip Prasad Tiwari

Kalinga University, Faculty of Pharmacy, Kotni, Naya Raipur, Chhattisgarh, India

Abstract--Melanoma is the most forceful sort of skin malignant growth and has exceptionally high paces of mortality. A beginning phase melanoma can be carefully eliminated, with an endurance pace of close to 100%. Notwithstanding, metastasized melanoma is challenging to fix. The 5-year endurance rates for patients with metastasized melanoma are still beneath 20%. Metastasized melanoma is presently treated by chemotherapy, designated treatment, immunotherapy and radiotherapy. The result of the majority of the ongoing treatments is not even close to hopeful. Despite the fact that melanoma patients with a transformation include the oncogene v-Raf murine sarcoma viral oncogene homolog B1 (BRAF) have an at first higher positive reaction rate to designated treatment, the greater part create obtained drug obstruction following a half year of the treatment. To increment treatment adequacy, early finding, more powerful pharmacological specialists, and more successful conveyance frameworks are critically required. Nanotechnology has been broadly read up for melanoma treatment and analysis, to diminish drug obstruction, increment helpful viability, and lessen secondary effects. Include this audit, we sum up the new advancement on the improvement of different nanoparticles for melanoma treatment and finding. A few normal nanoparticles, including liposome, polymerases, dendrites, carbon-based nanoparticles, and human egg whites, have been utilized to convey chemotherapeutic specialists, and little meddling ribonucleic acids (siRNAs) against flagging particles have additionally been tried for the treatment of melanoma. For sure, a few nanoparticle-conveyed drugs have been supported by the US Food and Drug Administration what's more, are at present include clinical preliminaries. The use of nanoparticles could deliver side results, which should be diminished so that nanoparticle-conveyed medications can be securely applied include the clinical setting.

Keywords---metastasis, early detection, nanoparticle-delivered, PI3K/Akt.

Introduction

Melanoma, started after melanocytes, is the most forceful kind appropriate to skin cancer. Despite the fact that melanoma reports just a tiny extent appropriate to skin disease frequency, it represents by far the most appropriate to skin disease deaths. Inasmuch as an instance, there were something else than 2 million new instances appropriate to skin disease include USA include 2012, also among them just 75,000 (3.75%) cases were melanoma. Be that as it may, melanoma represents 9000 (75%) passing's among complete 12,000 passing's brought about by skin malignant growth include 2012. beginning phase melanoma that consumes not blowout or metastasized towards different organs can be eliminated by medical procedure, with high endurance rates. It consumes been accounted inasmuch as that fix rate inasmuch as those patients with removable melanoma came towards 97%–99.8% subsequently chemosurgery. Be that as it may, metastasized irremovable melanoma should be treated by other standard treatments, like chemotherapy, radiotherapy, designated treatment, also immunotherapy. Unfortunately, treatment results are not good also patients' reactions to large portion appropriate to these treatments are very poor. As appropriate to now, most well-known drug utilized inasmuch as treatment appropriate to melanoma is decarbonize (DTIC), which is US Food and Drug Administration (FDA)- endorsed, first-line treatment inasmuch as patients with wild-type melanomas. Albeit include *in vitro* tests showed that DTIC created cytotoxicity include numerous melanoma cell lines, reaction towards DTIC was as it were 5%–10% when utilized alone. Utilization appropriate to designated treatment against changed oncogene v-Raf murine sarcoma biological oncogene homolog B1 (BRAF) showed an emphatically expanded reaction, coming towards 80%. Unfortunately, most patients will have obtained drug protection from BRAF inhibitors following half year appropriate to treatment, bringing about loss appropriate to clinical adequacy. middle endurance season appropriate to patients with metastasized melanoma is just 6–10 months, also 5-year endurance rate is under 20%.^{13–15} Therefore, gotten towards next level treatment productivity is direly required inasmuch as melanoma. Towards accomplish this, improvement include determination also treatment is required. Early location appropriate to melanoma can prompt careful expulsion before metastasis. ID appropriate to involved quality changes also comprehension appropriate to deregulations include key flagging pathways can help include plan appropriate to legitimate helpful systems. Effective treatment will likewise depend on disclosure appropriate to explicit, more strong pharmacological specialists furthermore, proficient conveyance frameworks. Nanotechnology includes plan, improvement, portrayal, also use appropriate to materials include Nano scale also, has been broadly read up inasmuch as clinical applications. Uses appropriate to nanotechnology include determination also treatment appropriate to melanoma have been concentrated broadly furthermore, have progressed investigation appropriate to melanoma infer aspects. Nanoparticles give powerful medication conveyance framework, permitting anticancer medications towards arrive at malignant growth site explicitly also include this way significantly further developing treatment

adequacy. Nanotechnology likewise has helped include analysis, by permitting imaging specialists towards be pressed into nanoparticles that span malignant growth. In this audit, we sum up new advancement include utilization appropriate to nanotechnology include conclusion also treatment appropriate to melanoma.

Normal nanoparticles utilized

In melanoma treatment Fast advancement include grasping science appropriate to disease cells has been made include previous ten years. Notwithstanding, treatment results appropriate to diseases have not changed lot. Disappointment appropriate to anticancer treatments is viewed as because appropriate to side impacts appropriate to most anticancer medications, low groupings appropriate to medications at growth site, also improvement appropriate to medication resistance. Nanoparticles have been concentrated widely inasmuch as drug conveyance, towards increment anticancer therapy adequacy. Contrasted also traditional medication conveyance draws near, nanoparticle-interceded conveyance appropriate to anticancer medications brings few wonderful benefits. Towards begin with, drugs conveyed by nanoparticles may have more drawn out organic life, due towards bundling security, also might be moved include site appropriate to disease because appropriate to upgraded penetrability also maintenance (EPR) at malignant growth sites. EPR is brought about by defectiveness appropriate to cancer vasculature as well as poor lymphatic drainage. In this manner, nanotechnology increments treatment adequacy what's more, diminishes incidental effects. Inasmuch as instance, Doxil (Janssen Biotech, Horsham, PA, USA), which is doxorubicin (Dox) figured out include liposomes containing polyethylene glycol (Stake), has 100-fold longer dissemination half-life also sevenfold lower cardio cytotoxicity than has free Dox. Second, nanoparticle can contain numerous medications also accordingly will work with blend treatment. It is known that mix appropriate to chemotherapy with immunotherapy or designated treatment creates significantly more powerful treatment outcomes. Third, nanoparticles can safeguard drugs made appropriate to delicate little meddling ribonucleic corrosive (siRNA) or proteins, from biochemical corruption include human body. This is because appropriate to covertness like elements appropriate to nanoparticles. Fourth, nanoparticles produce consistent arrival appropriate to conveyed drugs. Moreover, nanoparticles can be intended towards have different capacities, inasmuch as example, focusing towards disease cells also delivering picture contrast. Many nanoparticles have been read up inasmuch as treatment appropriate to melanoma, including liposomes, dendrites, polymerases, living nanoparticles, inert nanoparticles, also protein-based nanoparticles (Table 1).

A liposome is circle comprising of lipid bilayer that contains watery center inasmuch as hydrophilic drugs. Hydrophobic medications can be contained between two layers. Liposome was created over long time back also was first nanoparticle utilized include medicine. Liposome is ready by sonicating lipid, with an interaction including emulsification. Homogeneous nanosized liposomes can be accomplished by sifting through 0.2 μM membrane. Furthermore unambiguous ligands against growth antigens can be appended towards liposome surface so that nanoparticles can target disease cells specifically. Liposomes have been utilized

towards convey chemotherapeutic medications, immunocytokines, also siRNA, towards increment treatment adequacy inasmuch as melanoma. Similar underlying nanoparticles have been planned, inasmuch example, cub some also noisome, which comprise appropriate to lipids

Table I Characteristics and side effects of common nanoparticles used in the study of melanoma treatment

Nanoparticle	Sample drugs delivered for melanoma	Clinical use	Characteristics	Side effects	Possible solutions
Liposomes	Cisplatin Vincristine Doxorubicin	Doxil	High biocompatibility High biodegradability High cell uptake rates Low drug encapsulation efficacy High cost	Hand-foot syndrome (skin rash, swelling, redness, pain) Toxicity to normal cells due to positive charge	Increase drug loading to reduce liposomes to be used Use of natural lipids
Dendrimers	Doxorubicin Cisplatin Paclitaxel Vincristine	Not yet available	High stability High solubility High drug encapsulation Uniform size Low biodegradation	Surface charge Accumulation in kidney and reticuloendothelium	Attach anionic polymer Control the size
Polymersomes	Doxorubicin Paclitaxel Cyclodextrin	Not yet available	High biodegradability Controlled release Not uniform size	Accumulation in kidney and reticuloendothelium	Control the size
Carbon-based nanoparticles	Doxorubicin	Not yet available	High biocompatibility High photostability Small size Not biodegradable	Inflammation ROS production	Adjust the size
Human albumin	Paclitaxel	Nab-paclitaxel (Abraxane®)	High solubility High stability Low toxicity	Side effects related with delivered drugs	Reduce dosage for a specific drug through combination therapies

The applications appropriate to nanotechnology include conventional therapies appropriate to melanoma

Nanotechnology has been applied towards generally regular treatments inasmuch as melanoma. It has been shown that Nano delivery appropriate to medications inasmuch as chemotherapy, designated treatment, immunotherapy, also photodynamic treatment has incredibly expanded treatment efficacy.

Chemotherapy

Chemotherapy is most widely recognized ordinary treatment inasmuch as melanoma. Chemotherapeutic specialists kill both harmful what's more, ordinary cells that are include division. Therefore, they are exceptionally harmful, also this

restricts measurement towards be utilized. Normal chemotherapeutic specialists utilized include melanoma are DTIC, vinblastine, also temozolomide. reaction charges, be that as it may, are extremely low. It has been accounted inasmuch as that reaction towards DTIC, vinblastine, also temozolomide was 7%, 9.5%, furthermore, 28%, separately include clinical preliminary with 64 patients. Nanoparticles have been utilized towards convey chemotherapeutic specialists also shown towards increment treatment adequacy also diminish incidental effects. Zhang et al showed that conveying chemotherapeutic specialist Dox by gold nanoparticles was very compelling against melanoma cell line.⁴⁸ Lo Prete et al applied cholesterol-rich Nano emulsion (named "LDE") towards convey topside include mouse model appropriate to melanoma. It diminished side impacts, expanding greatest endured portion fivefold, also expanded restraint appropriate to cancer development by concentrating topside at growth site (a fourfold higher focus include growth than with free etoposide). Ndinguri et al pressed Dox include nanoparticle with extra immunizer against CD44, towards explicitly target dangerous cells.⁵⁰ nanoparticle decreased growth size by 60% contrasted also untreated cancer. Catch PTX (Abraxane) has been endorsed by FDA inasmuch as therapy appropriate to bosom malignant growth. Hersh et al revealed Phase II clinical preliminary utilizing grab PTX (Abraxane) include both beforehand treated also untreated melanoma patients. reaction rates were 2.7% also 21.6%, individually. include review, 22% chemotherapy-innocent patients ceased treatment because appropriate to grade 3 or 4 neuropathy, alopecia, neutropenia, also exhaustion. Kottschade et al⁵¹ directed Phase II clinical preliminary utilizing seize PTX also carboplatin include cutting edge melanoma, include chemotherapy-innocent also 35 beforehand treated melanoma patients. reaction rate was 25.6% include chemotherapy naive associate also was 8.8% include beforehand treated accomplice. extreme after effects were neutropenia, thrombocytopenia, neurosensory issues, weariness, queasiness, also regurgitating. include another clinical preliminary, Kottschade et al⁵² showed vascular endothelial development factor (VEGF) counter acting agent expanded impact appropriate to seize PTX. Ott et al showed that blend of B cell lymphoma protein (Bcl)-2 antisense oligonucleotide, temozolomide, also catch PTX created reaction appropriate to 40.6%.⁵³ Comparative secondary effects towards that uncovered include Kottschade et al's review were likewise detailed.

Designated treatment

Intracellular flagging pathways assume vital part include carcinogenesis also guess appropriate to numerous tumors, including melanoma, also numerous inhibitors have been created inasmuch as therapy appropriate to malignant growth (Table 2). These pathways are actuated either by hereditary deformities or ecological factors. Obesity is typical illness that is described by deregulated energy balance also metabolic disturbance. It can increment disease occurrence also results include more unfortunate anticipation through initiation appropriate to different sign pathways, including mitogen activated protein kinase (MAPK), phosphoinositide 3-kinase (PI3K), also signal transducer also activator appropriate to record 3 (STAT3). MAPK pathway is significant pathway include melanoma. It has been concentrated on widely include melanoma; also numerous inhibitors have been created towards restrain this pathway inasmuch


as treatment. Enactment appropriate to MAPK pathway include melanoma has been demonstrated towards be brought about by initiation appropriate to changes include its parts; transformation rate is 50%–70% inasmuch as BRAF, 15%–30% inasmuch as NRAS, 2% inasmuch as KRAS also 1% inasmuch as HRAS. PI3K/protein kinase B (Akt) pathway can synergistically influence MAPK pathway towards advance improvement appropriate to melanoma. Flagging pathways appropriate to MAPK also PI3K/ Akt also their crosstalk are displayed include Figure 1. two pathways direct an expansive scope appropriate to target proteins. 58,65 typical endurance pathway downstream appropriate to both MAPK also PI3K/Akt is mitochondrial apoptosis signal pathway that incorporates Bcl-2 group appropriate to proteins, including both proapoptotic furthermore, antiapoptotic signals. Moreover, hindrance appropriate to both MAPK also PI3K/Akt pathways has been displayed towards have synergistic impact include treatment appropriate to melanoma. Nanotechnology has been utilized towards convey specific inhibitors appropriate to MAPK pathway. Chitosan nanoparticle was utilized towards convey downstream objective protein VEGF siRNA also was shown towards further develop helpful effect. Yin et al utilized useful graphene oxide towards convey plasmid-based STAT3 siRNA also showed essentially diminished xenografted cancer growth. Indeed, STAT3 is considered as critical middle person include melanoma which advances cerebrum metastasis. Inhibition appropriate to phosphorylated STAT3 has been displayed towards increment adequacy appropriate to cancer corruption factor (TNF)- alpha inasmuch as melanoma. Nanoparticle has likewise been intended towards convey siRNA against oncogene c-Myc towards target melanoma cells B16F10 also exhibited adequacy against melanoma. Tran et al arranged nanoparticle towards contain both siRNAs against BRAF also, Akt3 which extraordinarily expanded anticancer impact

Table 2 Common inhibitors of the pMAPK and PI3K/Akt pathways

Signaling molecule	Inhibitors	References
RTK	Imatinib, dasatinib, sorafenib	Inamdar ⁵⁵
Ras	Tipifarnib, R115777, BMS-214662, L-778123	Inamdar ⁵⁵
Raf	Sorafenib, PLX4032, XL281, RAF-265	Inamdar ⁵⁵
Mek	AZD6244, U0126, PD0325901, CI-1040, XL518, AZD8330, ARRY-162, ARRY-300, perifosine, erucyl-phosphocholine	Chen ⁵⁴ Inamdar ⁵⁵
ERK	DHMEQ	Inamdar ⁵⁵
Akt	Ly294002, Bez235, PI103	Inamdar ^{54,55}
mTOR	Rapamycin, Bez235, PI103	Inamdar ^{54,55}
NF-kB	Curcumin, plumbagin	Inamdar ⁵⁵

Targeting mitochondrial pathway

As examined over, significant downstream middle person appropriate to PI3K/Akt also MAPK is mitochondrial apoptotic pathway. pathway is directed by proapoptotic proteins also antiapoptotic proteins. previous incorporates Bak, Bax, Bid, Bim, Bcl-related passing advertiser (BAD), Noxa, also Jaguar. last option incorporates Bcl-2, Bcl-xL, Bcl-w, Mcl-1, also, A1. Both PI3 K also MAPK can increment antiapoptotic protein articulation also decline proapoptotic protein articulation also consequently increment proportion appropriate to antiapoptotic proteins/proapoptotic proteins. This assumes critical part include melanoma commencement, support, also medication resistance. Inversion appropriate to this proportion, either by restraint appropriate to antiapoptotic proteins or overexpressing proapoptotic proteins, has been utilized inasmuch as treatment appropriate to melanoma. Inasmuch as instance, restraint appropriate to Bcl-2 by ABT-737 caused melanoma cell apoptosis.



The application appropriate to nanotechnology inasmuch as diagnosis appropriate to melanoma

The point appropriate to starting conclusion is towards make judgment about whether melanoma is available also at what formative stage is melanoma. An early finding is basic towards acquire high endurance rates inasmuch as melanoma patients. Past unaided eye assessment also histological perception, numerous new advancements have been utilized towards help conclusion, inasmuch as example, ceroscopy, all out body photography, reflectance confocal microscopy, multispectral computerized imaging investigation, also RNA microarray. include metastasized melanoma, sub-atomic marks appropriate to melanoma might be distinguished so that an appropriate treatment system can be made. This is significant, as sub-atomic changes are related with intracellular flagging pathway changes, clinical highlights, also reactions towards various treatment systems. Sub-atomic determination could be remembered inasmuch as normal determination inasmuch as melanoma include close future. More also all more recently evolved methods are utilized include atomic determination appropriate to melanoma, inasmuch as example, tissue cluster, proteomics, also DNA sequencing. During treatment appropriate to melanoma, judgment appropriate to adequacy is expected towards evaluate whether methodology utilized is fitting also whether there is any better method inasmuch

as expanding patient's reaction towards treatment. This assurance incorporates imaging growth size also, vascular design. Nanotechnology has been demonstrated towards besignificant instrument towards work on determination appropriate to melanoma. Nanoparticle quantum dabs (QDs) have been utilized inasmuch as early discovery appropriate to melanoma, due towards their photophysical properties, including excellent brilliance, size-tunable emanation frequency, wide excitation range with thin emanation profile, also astounding photostability.105,106 QDs can be formed with disease explicit atoms, inasmuch as example, folic corrosive or antibodies perceiving antigens overexpressed include malignant growth cells.107 Zheng et al utilized PEG-COOH covered profoundly fluorescent CdSe/ZnS center/shell QDs formed with immunizer against CD146 towards recognize CD146 overexpressed melanoma cells include refined also fixed cells.108 Flow cytometry furthermore, confocal microscopy showed positive cells towards have high splendor, photostability, also particularity. Kim et al fostered coculture framework containing both melanoma cells also typical melanocytes towards test QDs formed with antibodies (ab732 or Ab733) against melanoma. It was demonstrated way that melanoma cells could be recognized from melanocytes.

Theranostic Nano medicine include melanomaAs appropriate to late, theranostic Nano medicine has been created, which utilizes nanoparticles inasmuch as both determination also treatment. By utilizing limit appropriate to nanoparticles towards convey different specialists include solitary nanoparticle, imaging specialists are integrated into nanoparticles with goal that activity appropriate to remedial specialists can be observed include continuous design. This could assist with evaluating drug dissemination also delivery or towards assess drug reaction also viability. Data can likewise be valuable inasmuch as analysts, towards direct definition appropriate to new nanoparticles with wanted conveyance attributes. At present most generally utilized nanoparticles inasmuch as theranostic design are liposomes also polymerases. Imaging specialists inasmuch as attractive reverberation imaging (MRI), radionuclide imaging, furthermore, fluorescence imaging have been contemplated. Theranostic Nano medicine has previously been applied include melanoma study. Vannucci et al built nanoparticle include light appropriate to weighty chain appropriate to human protein ferritin towards test include melanoma.

Conclusion

Metastatic melanoma is as yet troublesome sickness towards treat. key inasmuch as further developing treatment viability is towards have better comprehension appropriate to pathogenesis appropriate to sickness, early analysis also recognizable proof appropriate to individual atomic composing, as well as novel also compelling medication conveyance frameworks. Nanotechnology has offered possibility towards work on conclusion also treatment appropriate to melanoma. It has been shown that nanoparticle-conveyed anticancer medications gather include growth tissue also hence increment treatment viability also diminish incidental effects. Nanoparticles have moreover been displayed towards convey imaging specialists towards recognize melanoma what's more, BRAF change. Further investigations are expected towards keep away from after effects brought about by some nanoparticles with goal that they can be applied clinically.

Reference

- Siegel R, DeSantis C, Virgo K, et al. Cancer treatment and survivorship statistics, 2012. *CA Cancer J Clin.* 2012;62(4):220–241.
- Siegel R, Naishadham D, Jemal A. Cancer statistics, 2012. *CA Cancer J Clin.* 2012;62(1):10–29.
- Jerant AF, Johnson JT, Sheridan CD, Caffrey TJ. Early detection and treatment of skin cancer. *Am Fam Physician.* 2000;62(2):357–368.
- American Cancer Society [homepage on the Internet]. Skin cancer facts. Available from: <http://www.cancer.org/cancer/cancercauses/sunanduvexposure/skin-cancer-facts>. Accessed June 5, 2013.
- Zitelli JA, Mohs FE, Larson P, Snow S. Mohs micrographic surgery for melanoma. *DermatolClin.* 1989;7(4):833–843.
- Boyle GM. Therapy for metastatic melanoma: an overview and update. *Expert Rev Anticancer Ther.* 2011;11(5):725–737.
- Katipamula R, Markovic SN. Emerging therapies for melanoma. *Expert Rev Anticancer Ther.* 2008;8(4):553–560.
- Khan KH, Goody RB, Hameed H, Jalil A, Coyle VM, McAleer JJ. Metastatic melanoma: a regional review and future directions. *Tumori.* 2012;98(5):575–580.
- Eigentler TK, Caroli UM, Radny P, Garbe C. Palliative therapy of disseminated malignant melanoma: a systematic review of 41 randomised clinical trials. *LancetOncol.* 2003;4(12):748–759.
- Serrone L, Zeuli M, Sega FM, Cognetti F. Dacarbazine-based chemotherapy for metastatic melanoma: thirty-year experience overview. *J ExpClin Cancer Res.* 2000;19(1):21–34.
- Brose MS, Volpe P, Feldman M, et al. BRAF and RAS mutations in human lung cancer and melanoma. *Cancer Res.* 2002;62(23):6997–7000.
- Davies H, Bignell GR, Cox C, et al. Mutations of the BRAF gene in human cancer. *Nature.* 2002;417(6892):949–954.
- Falkson CI, Ibrahim J, Kirkwood JM, Coates AS, Atkins MB, Blum RH. Phase III trial of dacarbazine versus dacarbazine with interferon alpha-2b versus dacarbazine with tamoxifen versus dacarbazine with interferon alpha-2b and tamoxifen in patients with metastatic malignant melanoma: an Eastern Cooperative Oncology Group study. *J ClinOncol.* 1998;16(5):1743–1751.
- Atkins MB, Hsu J, Lee S, et al; Eastern Cooperative Oncology Group. Phase III trial comparing concurrent biochemotherapy with cisplatin, vinblastine, dacarbazine, interleukin-2, and interferon alfa-2b with cisplatin, vinblastine, and dacarbazine alone in patients with metastatic malignant melanoma (E3695): a trial coordinated by the Eastern Cooperative Oncology Group. *J ClinOncol.* 2008;26(35):5748–5754.
- Sharma A, Sharma AK, Madhunapantula SV, et al. Targeting Akt3 signaling in malignant melanoma using isoselenocyanates. *Clin Cancer Res.* 2009;15(5):1674–1685.
- Bei D, Meng J, Youan BB. Engineering nanomedicines for improved melanoma therapy: progress and promises. *Nanomedicine(Lond).* 2010;5(9):1385–1399.
- Pacheco I, Buzea C, Tron V. Towards new therapeutic approaches for malignant melanoma. *Expert Rev Mol Med.* 2011;13:e33.

- Tran MA, Watts RJ, Robertson GP. Use of liposomes as drug delivery vehicles for treatment of melanoma. *Pigment Cell Melanoma Res.* 2009;22(4):388–399.
- Langer R. Drug delivery and targeting. *Nature.* 1998;392(6679 Suppl): S5–S10.
- Duncan R. The dawning era of polymer therapeutics. *Nat Rev Drug Discov.* 2003;2(5):347–360.
- Prabhakar U, Maeda H, Jain RK, et al. Challenges and key considerations of the enhanced permeability and retention effect for nanomedicine drug delivery in oncology. *Cancer Res.* 2013;73(8):2412–2417.
- Gabizon A, Catane R, Uziely B, et al. Prolonged circulation time and enhanced accumulation in malignant exudates of doxorubicin encapsulated in polyethylene-glycol coated liposomes. *Cancer Res.* 1994;54(4): 987–992.
- Moghimi SM, Hunter AC. Capture of stealth nanoparticles by the body's defences. *Crit Rev Ther Drug Carrier Syst.* 2001;18(6):527–550.
- Prabhu P, Patravale V. The upcoming field of theranostic nanomedicine: an overview. *J Biomed Nanotechnol.* 2012;8(6): 859–882.
- Sharifi S, Behzadi S, Laurent S, Forrest ML, Stroeve P, Mahmoudi M. Toxicity of nanomaterials. *ChemSoc Rev.* 2012;41(6):2323–2343.