

How to Cite:

Panigrahi, B., Dash, S., Nanda, S. S., & Satapathy, R. N. (2022). Study of association of maternal triglycerides level with neonatal macrosomia in non-obese non-diabetic pregnant women. *International Journal of Health Sciences*, 6(S2), 12317–12326.
<https://doi.org/10.53730/ijhs.v6nS2.8266>

Study of association of maternal triglycerides level with neonatal macrosomia in non-obese non-diabetic pregnant women

Bhagyashree Panigrahi

Senior Resident, Dept. Of O&G, PRM Medical College, Baripada, Odisha, India (Ex PG, SCB Medical College, Cuttack)

Subhalaxmi Dash

Asst. Professor, Dept. Of O&G, MKCG Medical College, Berhampur, Odisha, India

Sudhanshu Sekhara Nanda

Asst. Professor, Dept of O&G, PRM Medical College, Baripada, Odisha, India
Corresponding author email: sudhansu81@gmail.com

Rabi Narayan Satapathy

Asst Professor, Dept of O&G, SCB Medical College, Cuttack, Odisha, India

Abstract--Maternal dyslipidemia exerts potential and adverse impacts on pregnant women and newborns. However, the association between maternal serum triglycerides level and the risk of macrosomia has not yet been clearly studied. We did this study for association between the maternal serum triglycerides level and risk of macrosomia among the non-diabetic and non obese pregnant women. To find the associations of maternal triglyceride levels with neonatal macrosomia in non-obese and pregnant women without Diabetes Mellitus. SCB Medical College, Cuttack, Odisha, India. This was a descriptive cross-sectional study of 100 pregnant patients with BMI less than 30. The medical records of 100 non obese and non-diabetic pregnant women at term were prospectively analyzed. Maternal fasting serum triglyceride levels were measured during late pregnancy. Scatter diagram was used to analyze the variables associated with the risk of macrosomia. Maternal TG level was higher in patient with macrosomia babies (333.00 ± 51.62) compare to those without macrosomia (253.15 ± 81.02). This difference in mean was statistically significant (P value = 0.001) (Table 5). There was a positive correlation between birth weight and maternal triglyceride level which was statistically significant with correlation coefficient value of 0.278 (r) and p value of 0.005. Our findings highlight the importance of maternal lipid metabolism in fetal overgrowth and may have implications for the

etiology and primary prevention of macrosomia. However, further prospective investigations involving larger population and basic research studies are necessary to fully evaluate their clinical value and the mechanisms

Keywords---maternal triglyceride, macrosomia, non-diabetic pregnant women.

Introduction

Macrosomia is defined as a fetal birth weight equal to or greater than 4000g irrespective of gestational age, has a serious impact on maternal and fetal perinatal outcomes¹. For decades, the occurrence of macrosomia has been attributed to high maternal blood glucose. With regards to the Pedersen hypothesis, excess maternal glucose transfers through the placenta and stimulates islet cells and hyperinsulinemia, resulting in macrosomia². Maternal dyslipidemia exerts potential and adverse impacts on pregnant women and newborns. However, the association between maternal serum triglycerides level and the risk of macrosomia has not yet been clearly studied. We did this study for association between the maternal serum triglycerides level and risk of macrosomia among the non-diabetic and non obese pregnant women.

Macrosomia is associated with important neonatal complications such as birth trauma, hypoglycemia, and hematologic and respiratory complications³. Macrosomia is a major risk factor for obesity and type 2 diabetes mellitus in adolescence and young adulthood. Even though fetal macrosomia occurs more often in pregnancies of diabetic mothers, considerable numbers of infants with macrosomia are born from non-diabetic mothers (8–14%)⁴. Clausen et al in 2005 have assessed the risk of macrosomia in relation to early second trimester maternal serum lipids⁵; In the current study, we have focused on maternal serum lipids during the first weeks of the third trimester of pregnancy. This is because the possibility of developing a late impaired glucose metabolism after the screening time in some of the mothers could not be excluded.

There are also known risk factors for macrosomia, including maternal fasting glycemia, body mass index (BMI) prior to pregnancy, and HbA1c⁶. In the current study, we used healthy non-diabetic, non-obese and normotensive pregnant women to focus on the remaining risk factors after restricting our study to such normal mothers. Growth of the fetus depends on nutrients such as glucose, lipids, and amino acids.⁴ Therefore, in this study we evaluated the association between metabolic characteristics, especially serum triglyceride levels of the healthy non-diabetic and non-obese pregnant women in the first weeks of the third trimester of the pregnancy and its association with delivering macrosomic or large-for-gestational-age (LGA) neonates.

Maternal abnormal lipid profile can also induce excessive fetal growth⁷. The changes in lipid metabolism during pregnancy are characterized by fat accumulation in maternal depots during early pregnancy and development of hyperlipidemia later⁸. Recently, some maternal lipid parameters have been

identified as independent risk determinants of fetal overgrowth, especially in pregnancies complicated by GDM or with DM³⁷. Several studies have demonstrated that in non-diabetic pregnancies, birth weight and the risk of macrosomia or large for gestational age (LGA) were positively associated with maternal serum triglyceride (TG) levels⁹, and while other studies failed to find any association. Studies have shown that patterns of maternal dyslipidemia and the prevalence of macrosomia vary across populations and ethnic groups¹⁰. While a few studies have explored the relationship between maternal serum lipids and neonatal birth weight in Chinese population, their sample sizes were relatively small¹¹. In the present study, we sought to evaluate the association between maternal serum triglyceride levels at late gestation and the risk of macrosomia among a large group of women without DM.

The National Cholesterol Education Program has set guidelines for triglyceride levels¹²

Level (mg/dL)	Interpretation (mmol/L)	
< 150	< 1.70	Normal range – low risk
150–199	1.70–2.25	Slightly above normal
200–499	2.26–5.65	Some risk
500 or higher	> 5.65	Very high – high risk

The hypertriglyceridemia that occurs during pregnancy is due to both the increased production of triglyceride rich lipoproteins and the decreased clearance of triglyceride rich lipoproteins. The increased production of triglyceride rich lipoproteins by the liver is due to the increased lipolysis of triglycerides that occurs in adipocytes, which results in an increase in free fatty acids that are transported to the liver. In the liver, these free fatty acids are used in the synthesis of triglycerides, which are then packaged into VLDL and secreted by the liver. The high estrogen levels in the third trimester stimulate lipogenesis and VLDL production in the liver. The decrease in clearance of triglyceride rich lipoproteins is due to a decrease in lipoprotein lipase and hepatic lipase¹³. The decrease in hepatic lipase is due to the elevated estrogen levels, while the decrease in lipoprotein lipase is likely due to a combination of factors including insulin resistance and elevated estrogen levels. The triglyceride enrichment of LDL and HDL is due to an increase in CETP activity resulting in the transfer of triglyceride from VLDL to LDL and HDL and a decrease in hepatic lipase, which decreases the removal of triglycerides from these lipoprotein particles¹⁴.

Maternal dyslipidemia is linked to adverse perinatal outcomes. In animal studies, mice fed a hypercholesterolemic diet had increased rates of abortion, mortality, small litter sizes and lower birth weights of offspring. The offspring of hypercholesterolemic mothers also showed growth impairment and reduced renal function, effects only partially reversed by being fed a standard diet after delivery¹⁵ .. Variations in maternal lipid metabolism have an impact on fetal growth. In pregnancies of non-diabetic mothers, maternal triglyceride levels are correlated with fetal birth weight. Concentration of triglycerides in the third trimester is a stronger predictor of birth weight than glucose parameters¹⁶.

Maternal lipid metabolism is vital determinant of fetal development and growth. Maternal lipid profile at early and midpregnancy gestation is significantly associated with neonatal birth weight, and with the risk of macrosomia. Maternal serum TG does not directly cross the placenta, but the presence of lipoprotein receptors, various fatty acid-binding proteins, and lipase activity in the placenta enables the release of free fatty acids (FFAs) from TG and their efficient transport to the fetus. It is speculated that the increased hydrolysis of maternal TG by placental lipoprotein lipase to FFAs and the excessive delivery of fatty acids to the fetus may be partly responsible for the increased risk of macrosomia among women with hypertriglyceridemia.

Objective

To find the associations of maternal triglyceride levels with neonatal macrosomia in non-obese and pregnant women without DM.

Study Design

- Descriptive cross-sectional study of 100 pregnant patients with BMI less than 30.

Inclusion Criteria

- 1) Healthy pregnant women without diabetes and without obesity.
- 2) Pregnant mothers of BMI 18-29
- 3) live-born singleton pregnancy
- 4) delivery at 37–42 gestational weeks
- 5) naturally conceived
- 6) had integrated medical records and a clear gestational age.
- 7) Macrosomic babies.

Exclusion Criteria

- 1) Preterm labour prior to 37 weeks of gestational age.
- 2) Any abnormality or disorder in fetus or neonates.
- 3) pre-gestational DM or gestational diabetes mellitus (GDM) or type 2 DM
- 4) Multiple pregnancies
- 5) Pregnant mothers with BMI ≥ 30

Methods

In this study, the cases were taken from those who attended the antenatal clinics of the O&G department of SCBMCH, Cuttack or were admitted to the same department during the period October 2019 to September 2020. All the cases had antenatal check up in the antenatal period and they were advised for delivery and postpartum check up here. During the visit women were informed about the present study and the consent were taken. A detailed clinical assessment of the patient was performed in the outpatient department including history (diabetes, BMI & socioeconomic status, etc), general physical examination and obstetrical examination. Routine investigations were done during antenatal visits. Serum

triglyceride levels of non-obese and non-diabetic pregnant women who fulfilled inclusion criteria after 8-10 hours of fasting in late gestation. All women are generally healthy pregnant women carrying a single fetus having BMI between 18 kg per m² and 29 kg per m² without a history of diabetes prior to or in previous pregnancies and with a negative result from the diabetes screening test in the current pregnancy. Also took a blood sample of blood sugar level(GCT) to rule out PGDM/type2 DM/ GDM assessed 2hours after administering 75gms of oral glucose. The main outcome was measurement of neonatal birth weight. The medical records of 100 non obese and non-diabetic pregnant women at term were prospectively analyzed. Maternal fasting serum triglyceride levels were measured during late pregnancy. Scatter diagram was used to analyze the variables associated with the risk of macrosomia.

Statistical analysis

All the data collected were entered into the Microsoft Excel 2007 software and further analyzed in SPSS version 24 (IBM Inc. Chicago). All the categorical variables were expressed in terms of number/frequency and percentages. Association between two categorical variables were obtained by using Chi-squared test/Fischer Exact test. All the continuous variables were expressed in terms of mean and standard deviation. Significance level in comparison of means were obtained by independent sample t-test test. Histograms and box plots were used to describe the continuous variables. Bar and pie charts were used to describe the categorical variables. Correlation between two continuous variable was measured by person correlation coefficient and scatter plot was used to depict the correlation. A p value less than 0.05 was considered statistically significant.

Results

The current study aims to findout the associations of maternal triglyceride levels with neonatal macrosomia in a total of 100 non-diabetic and non-obese women. The mean age of the study participant was 26.21 ± 3.42 years which ranged from a minimum of 19 years to a maximum of 38 years. Mean weight gain during pregnancy was 11.08 ± 1.29 kilograms with a minimum of 8 kgs and a maximum of 14 kgs. Mean BMI of the study participants was 24.56 ± 2.03 Kg/m² . Mean 2-hour GCT after giving 75 grams of glucose was 100.81 ± 14.10 with a minimum of 75 and a maximum of 132 (Table-1)

Mean maternal triglyceride level was 254.5 ± 79.25 with a minimum of 100 to a maximum of 434 (Table 2). Mean birth weight was 3.85 ± 0.20 with a minimum of 3.6 to a maximum of 4.8. Similarly, RBS was 66.74 ± 15.60 with a minimum of 42 to a maximum of 99 (Table 3) Fetal outcome among the patient was given in table 4.. Macrosomia was present in 13% of the subjects . Total 22 neonates needed NICU admission. Neonatal complication like hypoglycemia and jaundice was seen in 11% each . Maternal TG level was higher in patient with macrosomia babies (333.00 ± 51.62) compare to those without macrosomia (253.15 ± 81.02). This difference in mean was statistically significant (P value = 0.001) (Table 5). There was a positive correlation between birth weight and maternal triglyceride level which was statistically significant with correlation coefficient value of 0.278 (r) and p value of 0.005 (figure 1). We did not find any statistical significant

difference with respect to BMI, RBS of newborn, weight gain in pregnancy, SBP and DBP.

Discussion

Here in our study, mean height of the subjects in the study was 156.39 ± 5.35 centimeters and the mean weight was 59.96 ± 5.03 kilograms. Mean weight gain during pregnancy was 11.08 ± 1.29 kilograms with a minimum of 8 kgs and a maximum of 14 kgs. Mean BMI of the study participants was 24.56 ± 2.03 Kg/m².

Blood sugar level & Blood pressure of the patients

Here in our study, mean 2-hour GCT after giving 75 grams of glucose was 100.81 ± 14.10 with a minimum of 75 and a maximum of 132 to exclude GDM/Type 2 DM. Mean systolic blood pressure was 112.52 ± 9.18 while the mean diastolic blood pressure was 74.38 ± 5.63 . On reviewing literature, association of maternal serum lipids at late gestation with the risk of neonatal macrosomia in women without diabetes mellitus by Xiangxiang Wang et al¹⁴⁵ in 2018, the mean SBP in mmHg was 120.07 ± 12.25 , mean DBP in mmHg was 76.63 ± 9.33 .

Triglyceride levels & newborn birth weight

Here the mean maternal triglyceride level was 254.5 ± 79.25 with a minimum of 100 to a maximum of 434. Mean birth weight was 3.85 ± 0.20 with a minimum of 3.6kg to a maximum of 4.8kg. Study of Association of maternal serum lipids at late gestation with the risk of neonatal macrosomia in women without diabetes mellitus by Xiangxiang Wang et al in 2018 showed the median of serum TG levels in the macrosomia group was much higher than that in the control group (3.52 mmol/L vs. 3.09 mmol/L, $p < 0.001$)¹⁷.

Prevalence of macrosomia

Here macrosomia was present in 13% of the subjects. Total 22 neonates needed NICU admission. Neonatal complication like hypoglycemia and jaundice was seen in 11% each. "Study of Association of maternal serum lipids at late gestation with the risk of neonatal macrosomia in women without diabetes mellitus" by Xiangxiang Wang et al in 2018 showed that the overall prevalence of macrosomia was 10.0%¹⁷.

Association of macrosomia with Maternal triglyceride levels

In our study maternal TG level was higher in patient with macrosomia babies (333.00 ± 51.62) compare to those without macrosomia (253.15 ± 81.02). This difference in mean was statistically significant (P value = 0.001). There was a positive correlation between birth weight and maternal triglyceride level which was statistically significant with correlation coefficient value of 0.278 (r) and p value of 0.005 (figure 1). We did not find any statistical significant difference with respect to BMI, RBS of newborn, weight gain in pregnancy, SBP and DBP. "Study of Association of maternal serum lipids at late gestation with the risk of neonatal

macrosomia in women without diabetes mellitus” by Xiangxiang Wang et al in 2018 showed that maternal serum triglyceride (TG) and highdensity lipoprotein cholesterol (HDL-C) levels were related to macrosomia; each 1 mmol/L increase in TG resulted in a 27% increase in macrosomia risk, while each 1 mmol/L increase in HDL-C level resulted in a 37% decrease in macrosomia risk, even after adjusting for potential confounders¹⁷.

Conclusion

Our results revealed that high maternal serum TG concentrations in late pregnancy and neonatal macrosomia are associated. Maternal TG level was higher in patient with macrosomia babies (333.00 ± 51.62) compare to those without macrosomia (253.15 ± 81.02). This difference in mean was statistically significant (P value = 0.001) . There was a positive correlation between birth weight and maternal triglyceride level which was statistically significant with correlation coefficient value of 0.278 (r) and p value of 0.005 . The combination of hypertriglyceridemia and low HDL-C levels was a stronger predictor of macrosomia than either alone. Our findings highlight the importance of maternal lipid metabolism in fetal overgrowth and may have implications for the etiology and primary prevention of macrosomia. However, further prospective investigations involving larger population and basic research studies are necessary to fully evaluate their clinical value and the mechanisms involved.

Table:1:Blood sugar level and blood pressure of the study participants

Blood sugar level (2hr GCT)(mg/dl)	Minimum	Maximum	Mean	SD
	75	132	100.81	14.10
Systolic BP(mmHg)	100	130	112.52	9.18
Diastolic BP(mmHg)	64	90	74.38	5.63

Table:2Maternal triglyceride level

	Minimum	Maximum	Mean	SD
Triglyceride level(mg/dl)	100	434	254.52	79.25

Table 3:Newborn birth weight and sugar level

	Minimum	Maximum	Mean	SD
Birth weight in Kg	3.60	4.80	3.85	0.20
RBS(mg/dl)	42	99	66.74	15.60

Table 4:Newborn outcome in the study population

Out comes		Number	Percentage
Macrosomia	Present	13	13
	Absent	87	87
NICU Admission	Needed	22	22
	Not needed	78	78
Neonatal complications	hypoglycemia	11	11
	Jundice	11	11
	None	78	78

Table 5:Association of macrosomia with Maternal triglyceride and other factors

Variables	Macrosomia present		Macrosomia absent		P value
	Mean	SD	Mean	SD	
Matrnl TG	253.15	81.02	333.00	51.62	0.001
BMI	24.67	2.00	23.81	2.13	0.926
RBS of newborn	67.57	15.24	61.15	17.47	0.654
SBP	101.95	14.21	93.15	10.85	0.347
DBP	112.18	9.09	114.77	9.88	0.878




Figure 1: Correlation between birthweight and maternal triglyceride level

Funding : None

Conflict of Interest: None

This study was approved by institutional ethics committee.

References

1. American Diabetes Association Diagnosis and classification of diabetes mellitus (position statement) *Diabetes Care*, 28 (Suppl. 1) (2005), pp. S37-S421
2. T. Harder, E. Rodekamp, K. Schellong, J.W. Dudenhausen, A. Plagemann Birth weight and subsequent risk of type 2 diabetes: a meta-analysis *Am J Epidemiol*, 165 (2007), pp. 849-857
3. J.M. Hawdon Babies born after diabetes in pregnancy: what are the short- and longterm risks and how can we minimise them? *Best Pract Res Clin Obstet Gynaecol*, 25 (2011), pp. 91-104
4. J. Pedersen Weight and length at birth of infants of diabetic mothers *Acta Endocrinol (Copenh)*, 16 (1954), pp. 330-342
5. G. Di Cianni, R. Miccoli, L. Volpe, C. Lencioni, A. Ghio, M.G. Giovannitti, et al. Maternal triglyceride levels and newborn weight in pregnant women with normal glucose tolerance *Diabet Med*, 22 (2005), pp. 21-25
6. Langer Fetal macrosomia: etiologic factors *Clin Obstet Gynecol*, 43 (2000), pp. 283- 297
7. Metzger BE, Lowe LP, Dyer AR, Trimble ER, Chaovarindr U, Coustan DR, Hadden DR, McCance DR, Hod M, McIntyre HD, et al. Hyperglycemia and adverse pregnancy outcomes. *N Engl J Med*. 2008;358:1991–2002. doi: 10.1056/NEJMoa0707943.
8. Vrijkotte TG, Algra SJ, Brouwer IA, van Eijsden M, Twickler MB. Maternal triglyceride levels during early pregnancy are associated with birth weight and postnatal growth. *J Pediatr* 2011; 159:736–742 e1
9. Son GH, Kwon JY, Kim YH, Park YW. Maternal serum triglycerides as predictive factors for large-for-gestational age newborns in women with gestational diabetes mellitus. *Acta Obstet Gynecol Scand*. 2010;89:700–704. doi: 10.3109/00016341003605677.
10. Retnakaran R, Ye C, Hanley AJ, Connelly PW, Sermer M, Zinman B, Hamilton JK. Effect of maternal weight, adipokines, glucose intolerance and lipids on infant birth weight among women without gestational diabetes mellitus. *CMAJ*. 2012;184:1353– 1360. doi: 10.1503/cmaj.111154.
11. Kitajima M, Oka S, Yasuhi I, Fukuda M, Rii Y, Ishimaru T. Maternal serum triglyceride at 24--32 weeks' gestation and newborn weight in nondiabetic women with positive diabetic screens. *Obstet Gynecol*. 2001;97:776–780. doi: 10.1016/S0029-7844(01)01328-X.
12. "Triglycerides". *MedlinePlus*. Archived from the original on 26 October 2012. Retrieved 2015-04-23.
13. Applebaum DM, Goldberg AP, Pykälistö OJ, Brunzell JD, Hazzard WR. Effect of estrogen on post-heparin lipolytic activity. Selective decline in hepatic triglyceride lipase. *J Clin Invest*. 1997 Apr; 59(4):601-8.
14. Reichen J, Karlaganis G, Kern F Jr. Cholesterol synthesis in the perfused liver of pregnant hamsters. *J Lipid Res*. 1987 Sep; 28(9):1046-52.

15. Kitajima M, Oka S, Yasuhi I, et al. Maternal serum triglyceride at 24--32 weeks' gestation and newborn weight in nondiabetic women with positive diabetic screens. *Obstet Gynecol.* 2001;97(5 Pt 1):776-780.
16. Nolan CJ, Riley SF, Sheedy MT, et al. Maternal serum triglyceride, glucose tolerance, and neonatal birth weight ratio in pregnancy. *Diabetes Care.* 1995;18(12):1550-1556.
17. Xiangxiang Wang, 1,2,3 Qingbo Guan, 1,2,3 Jiajun Zhao, 1,2,3 Feifei Yang, 1,2,3 Zhongshang Yuan, 4 Yongchao Yin, 5 Rui Fang, 1,2,3 Lingwei Liu, 1,2,3 Changting Zuo, 6 and Ling Gao 3,7 Association of maternal serum lipids at late gestation with the risk of neonatal macrosomia in women without diabetes mellitus