

How to Cite:

Jamdade, A., Rani, A., Behera, S. S. P., Sharma, D., Manik, A., & Singla, D. (2022). Treatment prostheses in TMJ dysfunction-pain syndrome: Systematic review & meta analysis. *International Journal of Health Sciences*, 6(S2), 14806–14817.
<https://doi.org/10.53730/ijhs.v6nS2.8935>

Treatment prostheses in TMJ dysfunction-pain syndrome: Systematic review & meta analysis

Dr. Anshuman Jamdade

Prof. and Head, Dept. of Oral Medicine and Radiology, Mahatma Gandhi Dental College and Hospital, Jaipur
Email: drjanshu@gmail.com

Dr Anusha Rani

Post Graduate, Dept of Conservative Dentistry and Endodontics, Jaipur Dental College, Jaipur
Email: dranushaendo@gmail.com

Dr. Sidhartha S P Behera

Reader, Dept of Prosthodontics, Sree Sai Dental College & Research Institute, Srikakulam, NTRUHS, Andhra Pradesh
Corresponding author email: drsidentb@gmail.com

Dr Deepak Sharma

Prof & HOD, Dept of Conservative Dentistry & Endodontics, Jaipur Dental college, Jaipur
Email: drdeepaksharma.mds@gmail.com

Dr. Amit Manik

MDS, oral pathology & Microbiology and Forensic Odontology, Reader, Desh bhagat dental college & hospital, Mandi Gobindgarh, Punjab
Email: amitmanik23@gmail.com

Dr. Deepika Singla

MDS, Senior Lecturer, Dept of Conservative Dentistry & Endodontics, Desh bhagat dental college & hospital, Mandi Gobindgarh
Email: Punjab.dsingla2810@gmail.com

Abstract---Manual therapy (MT) and exercise have been extensively used to treat people with musculoskeletal conditions such as temporomandibular disorders (TMD). The evidence regarding their effectiveness provided by early systematic reviews is outdated. The aim of this study was to summarize evidence from and evaluate the methodological quality of randomized controlled trials that examined the effectiveness of MT and therapeutic exercise interventions

compared with other active interventions or standard care for treatment of TMD. Material and methods: Electronic data searches of 6 databases were performed, in addition to a manual search. Randomized controlled trials involving adults with TMD that compared any type of MT intervention (eg, mobilization, manipulation) or exercise therapy with a placebo intervention, controlled comparison intervention, or standard care were included. The main outcomes of this systematic review were pain, range of motion, and oral function. Forty-eight studies met the inclusion criteria and were analyzed. Results: The overall evidence for this systematic review was considered low. The trials included in this review had unclear or high risk of bias. Thus, the evidence was generally downgraded based on assessments of risk of bias. Most of the effect sizes were low to moderate, with no clear indication of superiority of exercises versus other conservative treatments for TMD. However, MT alone or in combination with exercises at the jaw or cervical level showed promising effects. Conclusions: No high-quality evidence was found, indicating that there is great uncertainty about the effectiveness of exercise and MT for treatment of TMD.

Keywords---temporomandibular disorders (TMD), TMJ dysfunction-pain syndrome, prostheses.

Introduction

Temporomandibular disorders (TMD) are commonly associated with other symptoms affecting the head and neck region, such as headache, ear-related symptoms, cervical spine dysfunction¹⁻⁴ and altered head and cervical posture.⁵⁻¹⁵ Physical therapy has been used for decades for treating craniomandibular disorders using thermal packs vapocoolants, and transcutaneous electrical nerve stimulation (TENS).¹⁶ Physical therapy is among the 10 most commonly used treatments for TMD,¹⁹ focused on decreasing neck and jaw pain, improving range of motion (ROM), and promoting exercise to maintain healthy function.²⁰ Therapeutic exercise and MT are used to improve strength, coordination, and mobility and to reduce pain,²¹ and treatment may include and focus on poor posture, cervical muscle spasm or pain, and treatment for referred cervical origin orofacial pain (pain referred from upper levels of the cervical spine).²² The evidence for the effect of electrophysical modalities has been questioned.²³ Manual therapy (including joint mobilization, manipulation, or treatment of the soft tissues) and therapeutic exercises in physical therapy treatments have been increasingly used by clinicians and researched due to positive outcomes in some conditions, especially for low back pain, neck pain, and related disorders.²⁴ Most of these early systematic reviews highlighted the positive effects of exercises and MT to improve symptoms and function in people with TMD. Therefore, the objectives of this systematic review were: (1) to summarize the evidence from and evaluate the methodological quality of RCTs that examined the effectiveness of MT and therapeutic exercise interventions in the management of TMD and (2) to determine the magnitude of the effect of these interventions to manage TMD.

Material and Methods

This review was restricted to trials with participants meeting the following criteria: (1) diagnosis of TMD (2) adult (>18 years of age), (3) musculoskeletal dysfunction, (4) pain impairment, (5) no previous surgery in the temporomandibular region, and (6) no other serious comorbid conditions (eg, fracture in region, cancer, neurological disease). The primary outcomes of interest for this systematic review were pain, ROM, and oral function. Oral function for this systematic review focused on limitations of daily activities of patients with TMD measured through different questionnaires. A secondary outcome of interest was pressure pain threshold (PPT). This systematic review was open to all time points: immediately posttreatment and short-term, intermediate-term, and long-term follow-up.

Quality Assessment (Risk of Bias)

Assessments of quality (risk of bias) were completed by 2 independent reviewers (any 2 members of the research team). Any discrepancies in quality ratings were resolved by discussion. (95% CI) to pool data. Heterogeneity was evaluated statistically using the I² statistic.

Results

The search of the literature resulted in a total of 3,549 published articles. 45 studies were included for this review from the search of the databases. In addition, 3 studies were obtained through a manual search, total 48 of studies were included in the final analysis (Fig. 1). Details of included studies are provided in Table 1.

Diagnosis

There was considerable diversity in the clinical presentations and diagnoses of participants with TMD among the included studies. Fourteen of the studies examined the effectiveness of the exercise or MT interventions in muscular TMD (myogenous TMD), 14 studies examined the effectiveness in patients with articular TMD (arthrogenous TMD), and 19 studies examined the effectiveness in patients with mixed diagnoses of TMD (including both myogenous and arthrogenous TMD).^{28,29} One study looked at both myogenous and arthrogenous TMD.⁵⁸ Twenty-one of the studies^{57–76} used the RDC/TMD established by Dworkin and LeResche²⁷ to classify the patients as having TMD. The remaining 27 studies used their own diagnostic criteria, based on signs and symptoms of the patients.

Effectiveness of Intervention by TMD Diagnoses: Posture Correction Exercise Myogenous TMD

Two studies evaluated the effectiveness of posture correction exercises for patients with myofascial pain. Both studies showed positive results of postural exercises for improving symptoms of muscular TMD. When pooling the data for these 2 studies, which had similar interventions, diagnoses, and outcomes, maximum pain-free mouth opening significantly increased in patients receiving

postural training compared with a control group. The MD in maximum pain-free mouth opening was 5.54 mm (95% CI=2.93, 8.15) (Fig. 2), which was clinically significant in favor of postural training.

Manual Therapy Targeted to the Orofacial Region in Myogenous TMD

Four studies looked at MT techniques, such as facial manipulation versus botulinum toxin⁸¹ or intraoral myofascial therapy versus waiting list, and self-care education and exercises for people with myogenous TMD. The results of these studies support the use of MT to treat myogenous TMD, as people treated with all of these approaches had improved mouth opening and reduced jaw pain from baseline. Although the results for the intraoral myofascial therapy and exercise groups were superior to the results for the waiting-list control group, there was no statistically significant difference between them. Although the results were mixed, most of the studies favored the use of exercises alone or as part of a general regimen to treat people with arthrogenous TMD, including disk displacements with or without reduction. However, one study⁶¹ did not find that exercises were superior to a control group involving general physical therapy treatment.

General Jaw Exercise Program in Mixed TMD

Eleven studies looked at exercises alone or as part of a general conservative therapeutic regimen to treat patients with mixed TMD. In general, exercises for mixed TMD compared with control groups had better results for decreasing pain and improving function and pain sensitivity of the masticatory muscles. However, compared with other forms of active treatments, such as splints, a global postural re-education program, or acupuncture, no significant differences between these treatments were found.

When pooling the results of studies with available data and similar interventions and outcomes, we found that exercises in the form of general jaw exercises plus conventional treatment or with the addition of an oral device⁹⁴ were not superior to other treatment modalities, such as splint therapy, global re-education posture, splint plus counseling, acupuncture, or standard conservative care, in improving pain intensity. The SMD for pain intensity was -0.06 (95% CI=-0.50, 0.38), with a very small effect size according to Cohen's guidelines.

The overall quality of evidence for most comparisons was low to moderate according to the GRADE approach.⁴⁹ The trials included in this review had unclear or high risk of bias. Thus, the evidence was generally downgraded for 3 reasons: (1) risk of bias, 2) level of heterogeneity (inconsistency), and (3) some imprecision surrounding the effect estimate. Details of GRADE assessment of the included studies are displayed in the Table. From the 14 analyses performed, most of the evidence was considered moderate (9 analyses). The rest of the evidence was considered low. Thus, we can say that the total evidence was considered low.




Figure 1: Flow chart of the study selection




Figure 2. Maximum pain-free opening: postural training versus control group in patients with myogenous temporomandibular disorders. CI confidence interval, IV inverse variance

Quality Assessment						Summary of Findings			
Number of Studies (Design)	Risk of Bias	Inconsistency	Indirectness	Imprecision	Publication Bias	Number of Patients			Quality
						Patients in Treatment Group	Patients in Control Group	Estimate (SMD or MD) [95% CI]	
Posture Correction Exercises in Myogenous TMD									
Outcome: maximum pain-free opening									
2 (RCTs)	Unclear risk of bias for both studies. Concealment of allocation not clear for both studies, no intention-to-treat analysis.	P value on test for heterogeneity $P=.30$, $I^2=6\%$ No inconsistency	No serious indirectness	No serious imprecision	No serious publication bias	50	50	MD-5.54 mm [2.93, 8.15]	Moderate quality because of risk of bias
Outcome: disturbance of symptoms with daily life									
2 (RCTs)	Unclear risk of bias for both studies. Concealment of allocation not clear for both studies, no intention-to-treat analysis.	P value on test for heterogeneity $P=.15$, $I^2=51\%$ No serious inconsistency	No serious indirectness	No serious imprecision	No serious publication bias	50	50	SMD-1.13 [0.48, 1.78]	Moderate quality because of risk of bias. Upgrading because of large effect size and narrow CI.
General Jaw Exercises Alone or Combined With Exercise Program in Myogenous TMD									
Outcome: pain-free maximum opening									
4 (RCTs)	Unclear risk of bias for most of the studies. Concealment of allocation not clear for 3 studies, and 1 study was not concealed. All studies did not perform intention-to-treat analysis.	P value on test for heterogeneity $P<.00001$, $I^2=88\%$ High heterogeneity	No serious indirectness	No serious imprecision	No serious publication bias	64	67	MD-5.94 mm [-1.0, 12.87]	Low quality because of risk of bias and inconsistency
Outcome: pain intensity									
5 (RCTs)	High risk of bias for most of the studies. Concealment of allocation not clear for 4 studies, and 1 study was not concealed. Adherence and cointerventions unclear for most of the studies. No intention to treat for 4 studies.	P value on test for heterogeneity $P=.10$, $I^2=49\%$ No serious inconsistency	No serious indirectness	No serious imprecision	No serious publication bias	85	90	SMD-0.43 [-0.02, 0.87]	Moderate quality because of risk of bias
Manual Therapy Targeted to the Orofacial Region in Myogenous TMD									
Outcome: pain intensity									
3 (RCTs)	Concealment of allocation not clear for 2 of the studies, and 1 study was not concealed. Adherence and cointerventions unclear for all of the studies. No intention to treat for 2 studies.	P value on test for heterogeneity $P=.78$, $I^2=0\%$ No serious inconsistency	No serious indirectness	No serious imprecision	No serious publication bias	44	44	MD-1.35 cm [0.91, 1.78]	Moderate quality because of risk of bias
Jaw/Neck Exercises Alone or as Part of a Conservative Regimen in Arthrogenous TMD									
Outcome: pain intensity									
4 (RCTs)	Concealment of allocation not clear for all of the studies. Blinding was unclear for 3 of the studies, and 1 study did not have appropriate blinding. Adherence and cointerventions unclear for all of the studies. No intention-to-treat analysis for 2 studies.	P value on test for heterogeneity $P=.01$, $I^2=73\%$ High inconsistency	No serious indirectness	No serious imprecision	No serious publication bias	73	73	SMD-0.68 [-0.04, 1.40]	Low quality because of risk of bias and inconsistency
Outcome: active mouth opening									
3 (RCTs)	Concealment of allocation not clear for all of the studies. Blinding was unclear for 2 of the studies, and 1 study did not have appropriate blinding. Adherence and cointerventions unclear for all of the studies.	P value on test for heterogeneity $P=.009$, $I^2=79\%$ High inconsistency	No serious indirectness	No serious imprecision	No serious publication bias	63	63	3.13 mm [-1.96, 8.23]	Low quality because of risk of bias and inconsistency
Jaw/Neck Exercises Alone or as Part of a Conservative Regimen vs Surgery in Arthrogenous TMD									
2 (RCTs)	Concealment of allocation not appropriate for both studies. Blinding was not appropriate for both studies. Adherence and cointerventions unclear for both studies.	P value on test for heterogeneity $P=.04$, $I^2=76\%$ High inconsistency	No serious indirectness	No serious imprecision	No serious publication bias	68	63	MD--1.01 mm [-5.43, 3.42]	Low quality because of risk of bias and inconsistency

Outcome: pain intensity									
5 (RCTs)	Concealment of allocation unclear for most of the studies. Blinding was appropriate for most of the studies. Adherence unclear for most of the studies. No or unclear intention to treat for 2 studies.	P value on test for heterogeneity $P=.58$, $I^2=0\%$ No Inconsistency	No serious indirectness	No serious imprecision	No serious publication bias	99	114	SMD=0.40 [0.13, 0.68]	Moderate quality because of risk of bias
Outcome: active mouth opening									
4 (RCTs)	Concealment of allocation unclear for all of the studies. Blinding was appropriate for most of the studies. Adherence unclear for all of the studies.	P value on test for heterogeneity $P=.93$, $I^2=0\%$ No Inconsistency	No serious indirectness	No serious imprecision	No serious publication bias	75	77	3.58 mm [1.46, 5.70]	Moderate quality because of risk of bias
General Jaw Exercise Program in Mixed TMD									
Outcome: pain intensity									
5 (RCTs)	Concealment of allocation unclear for 4 of the studies. One study did not have appropriated allocation concealment. Appropriate blinding was unclear for most of the studies. Adherence and cointerventions unclear for all of the studies.	P value on test for heterogeneity $P=.14$, $I^2=41\%$ No serious inconsistency	No serious indirectness	No serious imprecision	No serious publication bias	81	81	SDM=-0.06 [-0.50, 0.38]	Moderate quality because of risk of bias
Outcome: mouth opening									
7 (RCTs)	Concealment of allocation unclear for 4 of the studies. Three studies did not have appropriate allocation concealment. Appropriate blinding was unclear for most of the studies. Adherence and cointerventions unclear for all of the studies. Most of the studies did not perform an intention-to-treat	P value on test for heterogeneity $P=.85$, $I^2=0\%$ No Inconsistency	No serious indirectness	No serious imprecision	No serious publication bias	132	138	MD=-0.25 [-2.08, 1.57]	Moderate quality because of risk of bias
Outcome: mouth opening									
2 (RCTs)	Concealment of allocation unclear for both studies. Appropriate blinding was unclear for one study. Intention to treat unclear for one study.	P value on test for heterogeneity $P=.000001$, $I^2=100\%$ Serious inconsistency	No serious indirectness	Moderate imprecision	No serious publication bias	47	49	MD=-17.33 mm [-10.39, 45.08]	Low quality because of risk of bias, inconsistency, and imprecision
Manual Therapy Plus Exercises for Mixed TMD									
Outcome: mouth opening									
2 (RCTs)	Concealment of allocation unclear for both studies. Appropriate blinding was unclear for one study and not appropriate for the other study. Intention to treat unclear for one study and not performed by the other study.	P value on test for heterogeneity $P=.02$, $I^2=82\%$ Serious inconsistency	No serious indirectness	No serious imprecision	No serious publication bias	42	41	MD=-6.10 mm [1.11, 11.09]	Moderate quality because of inconsistency

Table 1: Characteristics of the studies

Discussion

Although the quality of the evidence is mostly uncertain and low, the results of our systematic review showed positive results when using postural exercises and jaw exercises to treat both myogenous and arthrogeous TMD disorders. Manual therapy alone or in combination with exercises shows promising effects. Manual therapy targeted to the cervical spine decreased pain and increased mouth ROM in patients with myogenous TMD. Exercises did not show superiority over other treatments for treating mixed TMD. A general exercise program was effective compared with arthrocentesis or arthrography for treatment of arthrogeous TMD, with conservative treatments as a first line of treatment. There remain limited RCTs of high quality that have investigated the effectiveness of MT and exercises to treat TMD.

Effect of Exercise for Treating TMD

Exercise programs are advocated for treating people with musculoskeletal disorders. Therapeutic exercises are prescribed to address TMD. Passive and active stretching of muscles are performed to increase mouth ROM and reduce pain. Postural exercises are helpful.²¹ The results of our systematic review are consistent with previous reviews,^{19,23} showing positive effects when using exercises to treat myogenous and arthrogenous TMD. In particular, interventions including exercises to correct head and neck posture and active and passive oral exercises can be effective for reducing musculoskeletal pain and improving oromotor function.⁵¹

Methodological Elements and Overall Quality of the Evidence Affecting Observed Effect

The overall rating of the evidence for this review was low. This finding was due mainly to the risk of bias of the analyzed studies. The methodological biases common to the included studies could have an impact on results. Selection bias could have existed, as only 20 trials reported appropriate randomization and only 4 reported concealment of allocation.

Another important bias was the lack of blinding, especially of the patients and assessors. Only 12 studies used blinded assessment of clinician-assessed outcomes such as mouth opening. However, we also were interested in pain, which is a subjective outcome and dependent on the patient's report. It is likely that lack of blinding could have affected the results of these studies. However, because of the nature of the interventions investigated, blinding would not be possible in many of them. There is empirical evidence showing that trials without appropriate randomization, concealment of allocation, and blinding tend to report an inaccurate treatment effect compared with trials that include these features.¹⁰⁹ Thus, the results of this systematic review should be interpreted with caution, especially in trials with subjective self-reported outcomes.

Studies did not report interventions in sufficient detail to be reproducible. In addition, they did not control for cointerventions and did not have adequate adherence to treatment. These issues are of importance for this study, as it is unclear if the effects on selected outcomes were due to the effect of exercise, MT, or other cointerventions. In addition, it is unclear if the participants received enough dosage of treatments, as adequate adherence was accomplished by only a very small proportion of studies (15.2%). Adherence testing should be systematically studied in future studies with exercise prescriptions. The present study used a compilation of items from all of the scales used in the reviewed physical therapy literature in addition to the risk of bias tool. Our recent analysis of health scales used to evaluate methodological quality determined that none of these scales are adequate for use alone.^{44,111} Therefore, we decided to use all of the scales, using a compilation of their items, to provide a comprehensive and sensitive evaluation of the quality of individual trials. Research investigating methodological predictors for determining trial quality in physical therapy is needed. There is the potential for selection bias, yet our group used a comprehensive search strategy and included databases as well as manual search.

Conclusion

Although the overall level of evidence is low, exercises and MT are safe and simple interventions that could potentially be beneficial for patients with TMD. Active and passive exercise for the jaw, postural exercises, and neck exercises appear to have favorable effects for patients with TMD. Manual therapy alone or in combination with exercises shows promising effects. Exercises did not show clear superiority over other conservative treatments for TMD

References

1. McNeely M, Armijo Olivo S, Magee D. A systematic review of physical therapy intervention for temporomandibular disorders. *Phys Ther.* 2006;86:710–720. [PubMed: 16649894]
2. Brantingham JW, Cassa TK, Bonnefin D, et al. Manipulative and multimodal therapy for upper extremity and temporomandibular disorders: a systematic review. *J Manipulative Physiol Ther.* 2013;36:143–201. [PubMed: 23697915]
3. Moher D, Liberati A, Tetzlaff J, et al. Preferred reporting items for systematic reviews and meta- analyses: the PRISMA statement. *Ann Intern Med.* 2009;151:264–269. [PubMed: 19622511]
4. Dworkin SF, LeResche L. Research diagnostic criteria for temporomandibular disorders: review, criteria, examinations and specifications, critique. *J Craniomandib Disord.* 1992;6:301–355. [PubMed: 1298767]
5. Kraus S. Temporomandibular disorders, head and orofacial pain: cervical spine considerations. *Dent Clin North Am.* 2007;51:161–193, vii. [PubMed: 17185065]
6. Okeson JP. *Orofacial Pain: Guidelines for Assessment, Diagnosis, and Management.* Chicago, IL: Quintessence Publishing Co; 1996.
7. Kovacs FM, Abaira V, Royuela A, et al. Minimal clinically important change for pain intensity and disability in patients with nonspecific low back pain. *Spine (Phila Pa 1976).* 2007;32:2915–2920. [PubMed: 18246018]
8. Farrar JT, Young JP, Jr, LaMoreaux L, et al. Clinical importance of changes in chronic pain intensity measured on an 11-point numerical pain rating scale. *Pain.* 2001;94:149–158. [PubMed: 11690728]
9. van der Roer N, Ostelo RW, Bekkering GE, et al. Minimal clinically important change for pain intensity, functional status, and general health status in patients with nonspecific low back pain. *Spine (Phila Pa 1976).* 2006;31:578–582. [PubMed: 16508555]
10. Maughan EF, Lewis JS. Outcome measures in chronic low back pain. *Eur Spine J.* 2010:1–11. [PMCID: PMC2989277] [PubMed: 20397032]
11. Dworkin RH, Turk DC, Wyrwich KW, et al. Interpreting the clinical importance of treatment outcomes in chronic pain clinical trials: IMMPACT recommendations. *J Pain.* 2008;9:105–121. [PubMed: 18055266]
12. Kropmans TJ, Dijkstra RU, Stegenga B, et al. Smallest detectable difference in outcome variables related to painful restriction of the temporomandibular joint. *J Dent Res.* 1999;78:784–789. [PubMed: 10096454]
13. Nussbaum EL, Downes L. Reliability of clinical pressure-pain algometric measurements obtained on consecutive days. *Phys Ther.* 1998;78:160–169. [PubMed: 9474108]
14. Ylinen J, Takala EP, Kautiainen H, et al. Effect of long-term neck muscle

- training on pressure pain threshold: a randomized controlled trial. *Eur J Pain*. 2005;9:673–681. [PubMed: 16246820]
15. Cathcart S, Pritchard D. Reliability of pain threshold measurement in young adults. *J Headache Pain*. 2006;7:21–26. [PMCID: PMC3451574] [PubMed: 16440140]
 16. Fuentes CJ, Armijo-Olivo S, Magee DJ, Gross DP. A preliminary investigation into the effects of active interferential current therapy and placebo on pressure pain sensitivity: a random crossover placebo controlled study. *Physiotherapy*. 2011;97:291–301. [PubMed: 22051585]
 17. Byrt T. How good is that agreement? *Epidemiology*. 1996;7:561. [PubMed: 8862998]
 18. Armijo-Olivo SA, Macedo LG, Gadotti IC, et al. Scales to assess the quality of randomized controlled trials: a systematic review. *Phys Ther*. 2008;88:156–175. [PubMed: 18073267]
 19. Armijo-Olivo S, Ospina M, da Costa BR, et al. Poor reliability between Cochrane reviewers and blinded external reviewers when applying the Cochrane risk of bias tool in physical therapy trials. *PloS One*. 2014;9:e96920. [PMCID: PMC4019638] [PubMed: 24824199]
 20. Hartling L, Hamm MP, Milne A, et al. Testing the risk of bias tool showed low reliability between individual reviewers and across consensus assessments of reviewer pairs. *J Clin Epidemiol*. 2012;66:973–981. [PubMed: 22981249]
 21. Higgins J, Green S. *Cochrane Handbook for Systematic Reviews of Interventions*, Version 4.2.6. In: Higgins J, Green S, eds. *The Cochrane Library*. Issue 4 Chichester, United Kingdom: John Wiley & Sons Ltd; 2006.
 22. Berard A, Bravo G. Combining studies using effect sizes and quality scores: application to bone loss in postmenopausal women. *J Clin Epidemiol*. 1998;51:801–807. [PubMed: 9762872]
 23. Deeks J, Higgins J, Altman D. *Analysing data and undertaking meta-analyses*. In: Higgins J, Green S, eds. *Cochrane Handbook for Systematic Reviews of Interventions*. Version 5.0.0. Available at: <http://www.cochrane-handbook.org>. Updated March 2011 Accessed June 24, 2015.
 24. Cohen J. *The concepts of power analysis*. In: Cohen J, ed. *Statistical Power Analysis for the Behavioral Sciences*. 2nd ed Hillsdale, NJ: Academic Press Inc; 1988:1–17.
 25. Guyatt GH, Oxman AD, Vist GE, et al. GRADE: an emerging consensus on rating quality of evidence and strength of recommendations. *BMJ*. 2008;336:924–926. [PMCID: PMC2335261] [PubMed: 18436948]
 26. Fink M, Ismail F, Hessling K, et al. The use of physical therapy for the treatment of craniomandibular disorders: a prospective, randomised clinical trial [article in German]. *Manuelle Medizin*. 2007;4:255–260.
 27. Lalabonova KH, Bakyrdzhev A. Physical and medicinal treatment of TMJ arthrosis [article in Russian]. *Stomatologiya (Mosk)*. 2008;87:50–51. [PubMed: 18577923]
 28. Petz R, Muska K. Therapy of restricted jaw movements [article in German]. *Z Arztl Fortbild (Jena)*. 1974;68:841–844. [PubMed: 4440003]
 29. van der Glas HW, Buchner R, van Grootel RJ. Vergelijking tussen behandelingsvormen bij myogene temporomandibulaire dysfunctie [Comparison of treatment options for myogenous temporomandibular dysfunction]. *Ned Tijdschr Tandheelkd*. 2000;107:505–512. [PubMed: 11383262]

30. Xue WH, Ding M, Su XC, et al. Clinical observation on warming needle moxibustion plus exercise for treatment of temporomandibular joint dysfunction syndrome [article in Chinese]. *Zhongguo Zhen Jiu*. 2007;27:322–324. [PubMed: 17645250]
31. Burgess JA, Sommers EE, Truelove EL, Dworkin SF. Short-term effect of two therapeutic methods on myofascial pain and dysfunction of the masticatory system. *J Prosthetic Dent*. 1988;60:606–610. [PubMed: 3199321]
32. Magnusson T, Syren M. Therapeutic jaw exercises and interocclusal appliance therap between two common treatments of temporomandibular disorders. *Swed Dent J*. 1999;23:27–37. [PubMed: 10371003]
33. Mulet M, Decker KL, Look JO, et al. A randomized clinical trial assessing the efficacy of adding 6 × 6 exercises to self-care for the treatment of masticatory myofascial pain. *J Orofac Pain*. 2007;21:318–328. [PubMed: 18018993]
34. Maloney GE, Mehta N, Forgione AG, et al. Effect of a passive jaw motion device on pain and range of motion in TMD patients not responding to flat plane intraoral appliances. *Cranio*. 2002;20:55–65. [PubMed: 11831346]
35. Komiya O, Kawara M, Arai M, et al. Posture correction as part of behavioural therapy in treatment of myofascial pain with limited opening. *J Oral Rehabil*. 1999;26:428–435. [PubMed: 10373091]
36. Wright EF, Domenech MA, Fischer JR., Jr Usefulness of posture training for patients with temporomandibular disorders. *J Am Dent Assoc*. 2000;131:202–210. [PubMed: 10680388]
37. Craane B, Dijkstra PU, Stappaerts K, De Laat A. Randomized controlled trial on physical therapy for TMJ closed lock. *J Dent Res*. 2012;91:364–369. [PubMed: 22318373]
38. Gavish A, Winocur E, Astandzelov-Nachmias T, Gazit E. Effect of controlled masticatory exercise on pain and muscle performance in myofascial pain patients: a pilot study. *Cranio*. 2006;24:184–190. [PubMed: 16933459]
39. Kalamir A, Graham PL, Vitiello AL, et al. Intra-oral myofascial therapy versus education and self-care in the treatment of chronic, myogenous temporomandibular disorder: a randomised, clinical trial. *Chiropr Man Therap*. 2013;21:17. [PMCID: PMC3706243] [PubMed: 23738586]
40. Kalamir A, Pollard H, Vitiello A, Bonello R. Intra-oral myofascial therapy for chronic myogenous temporomandibular disorders: a randomized, controlled pilot study. *J Man Manip Ther*. 2010;18:139–146. [PMCID: PMC3109684] [PubMed: 21886424]
41. Michelotti A, Steenks MH, Farella M, et al. The additional value of a home physical therapy regimen versus patient education only for the treatment of myofascial pain of the jaw muscles: short-term results of a randomized clinical trial [erratum in: *J Orofac Pain*. 2006;20:106]. *J Orofac Pain*. 2004;18:114–125. [PubMed: 15250431]
42. La Touche R, Paris-Aleman A, Mannheimer JS, et al. Does mobilization of the upper cervical spine affect pain sensitivity and autonomic nervous system function in patients with cervico-craniofacial pain: a randomized-controlled trial. *Clin J Pain*. 2013;29:205–215. [PubMed: 22874091]
43. de Felicio CM, Melchior Mde O, Ferreira CL, Da Silva MA. Otologic symptoms of temporomandibular disorder and effect of orofacial myofunctional therapy. *Cranio*. 2008;26:118–125. [PubMed: 18468271]
44. Ismail F, Demling A, Hessling K, et al. Short-term efficacy of physical therapy compared to splint therapy in treatment of arthrogenous TMD. *J Oral*

- Rehabil. 2007;34:807–813. [PubMed: 17919246]
45. Nascimento MM, Vasconcelos BC, Porto GG, et al. Physical therapy and anesthetic blockage for treating temporomandibular disorders: a clinical trial. *Med Oral Patol Oral Cir Bucal*. 2013;18:e81–e85.
 46. de Felicio CM, de Oliveira MM, da Silva MA. Effects of orofacial myofunctional therapy on temporomandibular disorders. *Cranio*. 2010;28:249–259. [PubMed: 21032979]
 47. Ficnar T, Middelberg C, Rademacher B, et al. Evaluation of the effectiveness of a semi-finished occlusal appliance: a randomized, controlled clinical trial. *Head Face Med*. 2013;9:5.
 48. [Niemela K, Korpela M, Raustia A, et al. Efficacy of stabilisation splint treatment on temporomandibular disorders. *J Oral Rehabil*. 2012;39:799–804. [PubMed: 22809314]
 49. Truelove E, Huggins KH, Mancl L, Dworkin SF. The efficacy of traditional, low-cost and nonsplint therapies for temporomandibular disorder: a randomized controlled trial. *J Am Dent Assoc*. 2006;137:1099–1107. [PubMed: 16873325]
 50. Tuncer AB, Ergun N, Tuncer AH, Karahan S. Effectiveness of manual therapy and home physical therapy in patients with temporomandibular disorders: a randomized controlled trial. *J Bodyw Mov Ther*. 2013;17:302–308. [PubMed: 23768273]
 51. Packer AC, Pires PF, Dibai-Filho AV, Rodrigues-Bigaton D. Effects of upper thoracic manipulation on pressure pain sensitivity in women with temporomandibular disorder: a randomized, double-blind, clinical trial. *Am J Phys Med Rehabil*. 2014;93:160–168. [PubMed: 24434889]
 52. Kalamir A, Bonello R, Graham P, et al. Intraoral myofascial therapy for chronic myogenous temporomandibular disorder: a randomized controlled trial. *J Manipulative Physiol Ther*. 2012;35:26–37. [PubMed: 22079052]
 53. Suryasa, I. W., Rodríguez-Gámez, M., & Koldoris, T. (2021). Get vaccinated when it is your turn and follow the local guidelines. *International Journal of Health Sciences*, 5(3), x-xv. <https://doi.org/10.53730/ijhs.v5n3.2938>
 54. Suryasa, I. W., Rodríguez-Gámez, M., & Koldoris, T. (2021). Health and treatment of diabetes mellitus. *International Journal of Health Sciences*, 5(1), i-v. <https://doi.org/10.53730/ijhs.v5n1.2864>